arXiv:1202.4104v1 [math.PR] 18 Feb 2012

The K-process on a tree as a scaling limit of the GREM-like trap model

L. R. G. Fontes [*][†] R. J. Gava [‡][§] V. Gayrard [¶][‖]

Abstract

We introduce trap models on a finite volume k-level tree as a class of Markov jump processes with state space the leaves of that tree. They serve to describe the GREM-like trap model of Sasaki-Nemoto. Under suitable conditions on the parameters of the trap model, we establish its infinite volume limit, given by what we call a K process in an infinite k-level tree. From this we deduce that the K-process also is the scaling limit of the GREM-like trap model on *extreme* time scales under a *fine tuning* assumption on the volumes.

AMS 2010 Subject Classifications: 82C44,60K35,60G70

Keywords and Phrases: random dynamics, random environments, K-process, scaling limit, trap models

1 Introduction

The long time behavior of slow dynamics in random environments and phenomena like *aging* is a research theme of recent interest. Trap models and related stochastic processes have been proposed as simple models where these issues can be studied and understood on a rigorous basis. Perhaps the simplest such models are Markov jump processes on given graphs with simple symmetric random walks as embedded chains. The mean jump times at the vertices are random iid parameters, with heavy tailed distribution, that may be seen as the depths of traps, playing the role of the random environment. The case of $\mathbb{Z}^d$ was extensively analysed in the physics [1, 2] as well as mathematical literature [3, 4, 5, 6]. The case of the *complete graph* was introduced in [7] as a toy model for the aging behavior of the REM, and is well understood [7, 8, 9, 10, 11]. The actual REM dynamics (with Gibbs factors instead of the iid heavy tailed random variables) was studied in [12, 13], where it is shown that aging is the same as in the complete graph. Refined understanding of this dynamics on a wide range of

---

[*] IME-USP, Rua do Matão 1010, 05508-090 São Paulo SP, Brazil, lrenato@ime.usp.br

[†] Partially supported by CNPq grant 305760/2010-6, and FAPESP grant 2009/52379-8

[‡] IME-USP, Rua do Matão 1010, 05508-090 São Paulo SP, Brazil, gavamat@yahoo.com.br

[§] Supported by FAPESP fellowship 2008/00999-0

[¶] CMI, 39 rue Joliot Curie, 13453 Marseille, France, veronique@gayrard.net

[‖] Partially supported by FAPESP grants 2007/59096-6 and 2009/51609-0

time scales was obtained in [12, 13, 15, 16, 14, 17]. A natural next step to the analyses of the REM is to consider correlated Hamiltonians, namely the p-spin SK models and the GREM. The p-spin dynamics was studied in [18, 19, 20] in a particular range of time scales and temperature parameters where aging is the same as in the REM. At present however there is no rigorous results about the GREM dynamics. The only results available are non rigorous theoretical results [7, 21, 22] and concern trap-like models of this dynamics. In this work we consider one of these models, namely the GREM-like trap model introduced by Sasaki and Nemoto [21].

Let us first describe a model with a fixed deterministic environment, which we call the *trap model on a tree*, and come back to the GREM-like trap model after that. Let $M_1, \ldots, M_k$ be positive integers, and consider a k-level rooted tree whose first generation has size M_1, and such that each vertex in generation $j-1$ has M_j offspring at generation j, $j = 2, \ldots, k$. The state space of our model are the leaves of that tree. The parameters of the model are as follows. To each leaf vertex we will attach a positive parameter γ_k, dependent on the vertex. To interior vertices (not leaves), we attach probabilities p_j, $j = 1, \ldots, k-1$, also dependent on the vertex, all of them positive, except for p_{root}, which vanishes. See Figure 1 below. The transition mechanism of the trap model is as follows. Once in a given leaf vertex x of the tree, the process waits an exponential time with mean $\gamma_k(x)$ and then jumps. The destination of the jump, another leaf vertex of the tree, let us call it y, is chosen as follows. An ancestor of x on the tree is first chosen by going up the path from x to the root, and independently flipping coins whose probabilities of heads are the p_j's encountered along the way, until tails come up for the first time. The corresponding stopping vertex is the chosen ancestor, let us call it z. We then choose y uniformly among the leaf vertices descending from z. The trap model on a tree is thus fully described.

The GREM-like trap model is the trap model on a (k-level) tree for which the γ_k's as well as the inverse of the p_j's, $j = 1, \ldots, k-1$, are random variables, independent over the vertices, whose common distribution on a given level j is in the basin of attraction of a stable distribution with index α_j, and such that $0 < \alpha_j < 1$, $j = 1, \ldots, k$. We are interested in the long time behavior of that model as volume diverges.

There is of course an issue of how time scales with the volumes, and how the volumes relate to each other. In this paper we derive a scaling limit for the process at times of the order of the maxima of the γ_k's – we call this time scale *extreme* scale. The volumes will be taken related to each other in what we call the *fine tuning* regime (see (6.6) on Section 6). Aging does not take place on this time scale. It rather requires taking a double limit procedure, first sending the volume to infinity, and then the time to zero (as discusssed e.g. in [10]). This will be done in a follow-up paper. Shorter time scales and different choices of the volume are studied in [23].

The scaling limit for the GREM-like trap model is derived in Section 6 – see Theorem 6.2 – as a particular case of an infinite volume limit of the trap model on a tree described above. In order to perform the latter limit, we consider an alternative description of that dynamics, since the original one is not straightforwardly conducive to an infinite volume version. This is done in Section 3; this representation, a key element of the paper, readily suggests an infinite volume limit version of the finite volume dynamics. In Section 4 we introduce such an infinite volume dynamics, and in the following Section 5, we prove a convergence result of the finite volume process to the infinite volume one – see Theorem 5.1.

In the remainder of this introduction, we discuss the alternative description of the trap model on a tree, and the infinite volume process. It is based on an inductive construction which we now outline. The induction is in the number of levels of the tree. Let us start with the 1-level case. This is the case studied in [9], and we use the same construction as in that reference. For each leaf vertex, we take a Poisson point process of rate 1, independent of each other; each point of each process carries the label of the respective leaf vertex. The superposition of the labeled points of all those processes in a single time line gives us a realization of the embedded chain of the 1-level trap model on a tree (whose transitions consist of uniform jumps on $\{1, \dots, M_1\}$). In order to complete the picture, it remains to give the sojourn times at each point s, and this is given by $\gamma_1(x)T_s^{(1)}$, where x is the label of s, $T_s^{(1)}$ are mean one exponentials, independent of each other as s varies. This ends the description of the 1-level model. Given a construction of the l-level model, $l \geq 1$, call it X_l, in order to construct the $l+1$-level model, we consider the partition of the nonnegative line into constancy intervals of X_l; at this point we should remark that X_l is a jump process, and we admit jumps to the same point, and, along with that, consecutive (distinct) constancy intervals of X_l with the same value of the process. This gives a realization of the l first coordinates of the embedded chain of X_{l+1}. In order to get the $l+1$-st coordinate, a uniform random variable in $\{1, \dots, M_{l+1}\}$, we again consider Poisson point processes of rate 1, one for each possible value of that coordinate, independent of each other and of X_l, and superimpose them on the time line of X_l, each point carrying the label coming from the respective $l+1$-st coordinate value. To make sure that each constancy interval of X_l gets a point, we introduce extra points, one placed at the left endpoint of each constancy interval. Each point carries an $l+1$-coordinate label, the first l coordinates of which are given by the value of X_l at the respective constancy interval, and the $l+1$-st coordinate is given by the label of either the respective Poisson point, or an independent random variable uniformly distributed in $\{1, \dots, M_{l+1}\}$, in the case of an extra point. This gives us the embedded chain of X_{l+1}. And the sojourn time at a point s is given by $\gamma_{l+1}(x)T_s^{(l+1)}$, where x is the label of s, $T_s^{(l+1)}$ are mean one exponentials, independent of each other and X_l as s varies, similarly as for the 1-level case. We need of course a consistency between γ_l of the l-level model and p_l of the $l+1$-level one, and this is established by the condition $p_l = 1/(1 + M_{l+1}\gamma_l)$.

A key fact making the above representation work is as follows. In an attempt to represent the $l+1$-level dynamics from the l-level one, it is natural to consider the number of coin flips of the $l+1$-level dynamics up to the first heads (of the first coin it flips, namely the one with heads probability p_l). This is a geometric random variable starting at 1. In the inductive construction, this should correspond to the number of points inside a constancy interval of X_l. Since the length of such an interval is exponentially distributed, we have that the number of (independent) Poisson points inside is geometric starting at 0, and the extra point yields a match on the form of the distribution, and the consistency condition matches the parameters.

The construction of the infinite volume dynamics is the natural analogue of the above finite volume one. The difference is that we have an infinite number of Poisson processes at each inductive step – no extra points now – so that there is a dense collection of points at each stage to attach sojourn times to, which we do in the same way as for finite volume. To have things make sense in the end, we of course need to have the sojourn times in any bounded interval summable, and that imposes a condition on the γ_l's. See Section 4, in particular (4.2). The limiting process is not a jump process, even though it may be shown to be Markovian. In order to make it càdlàg, we need to have it attain infinities at certain (exceptional) times (see (4.6) and Remark 4.7).

2 The model

We describe the trap model on a tree in detail now. Let us start with the tree. Throughout, k will be a fixed integer in $\mathbb{N}_* := \{1, 2, \dots\}$. Consider k numbers $M_1, \dots, M_k \in \mathbb{N}_*$, sometimes below called volumes, and let $\mathcal{M}_j = \{1, \dots, M_j\}$, $\mathcal{M}|_j = \mathcal{M}_1 \times \dots \times \mathcal{M}_j$, $j = 1, \dots, k$. Let us then consider the tree rooted at $\emptyset$

$$\mathbb{T}_k^F = \cup_{j=0}^k \mathcal{M}|_j, \tag{2.1}$$

where $\mathcal{M}|_0 = \{\emptyset\}$. We will use the notation $x|_j \equiv x_1 \dots x_j$ for a generic element of $\mathcal{M}_j$. We will also use the notation

$$\mathcal{M}|^j = \mathcal{M}_j \times \dots \times \mathcal{M}_k, \tag{2.2}$$

$1 \leq i \leq k$, and let $x|^j \equiv x_j \dots x_k$ denote a generic element of $\mathcal{M}|^j$.

$\mathbb{T}_k^F$ is of course a finite tree and this is emphasized in the notation by the use of the superscript "F". We understand the root to be at the 0-th generation of $\mathbb{T}_k^F$, and $x|_j \in \mathcal{M}|_j$ to be in its j-th generation, $1 \leq j \leq k$. For $0 \leq i < j \leq k$, we will regard $x|_i$ as an ancestor of $x|_j$, $j = 1, \dots, k$. We will sometimes use simply x for $x|_k$.

Let $\bar{\mathbb{N}}_* = \mathbb{N}_* \cup \{\infty\}$. In the sections below, we will consider the infinite tree

$$\mathbb{T}_k = \cup_{j=0}^k \bar{\mathbb{N}}_*^j, \tag{2.3}$$

where $\bar{\mathbb{N}}_*^0 = \{\emptyset\}$, with generations and ancestors as in $\mathbb{T}_k^F$.

The dynamics we will consider is a continuous time Markov jump process on the set of leaves of $\mathbb{T}_k^F$, namely its k-th generation $\mathcal{M}|_k$.

Let us describe the transition mechanism of the process. There will be a set of parameters for that. In order to distinguish this finite tree description from the later infinite tree one (to be presented in Section 4 below), on the one hand, and to emphasize the analogy between the two cases, on the other hand, we continue resorting to the use the superscript "F" for the set of parameters of the finite volume process as well.

For $j = 1, \dots, k-1$, let

$$p_j^F : \mathcal{M}|_j \to (0, 1), \tag{2.4}$$

and

$$\gamma_k^F : \mathcal{M}|_k \to (0, \infty). \tag{2.5}$$

For $x \in \mathcal{M}|_k$, let $g_x \in \{0, 1, \dots, k-1\}$ be a random variable such that

$$P(g_x = i) = [1 - p_i^F(x|_i)] \prod_{j=i+1}^{k-1} p_j^F(x|_j), \tag{2.6}$$

where by convention $\Pi_{j=k}^{k-1} p_j^F(x|_j) = 1$ and $p_0^F \equiv 0$.

Let Z_k^F be a continuous time Markov chain on $\mathcal{M}|_k$ as follows. When Z_k^F is at $x \in \mathcal{M}|_k$, it waits an exponential time of mean $\gamma_k^F(x)$ and then jumps as follows. It first looks at a copy g_x' of g_x (at each time independent of the copies looked at previously). If $g_x' = j$, then, letting $a_x(j)$ denote the (only) *ancestor* of x on generation j of $\mathbb{T}_k^F$ (namely $a_x(j) = x|_j$), Z_k^F jumps uniformly at random to one of the *descendants* of $a_x(j)$ in $\mathcal{M}|_k$. In other words, given that $g_x' = j$, then the coordinates $x|_j (= a_x(j))$ of x are left unchanged, and the remainder coordinates are chosen uniformly at random on $\mathcal{M}|^{j+1}$. We may then say that the transition distribution of the jump



Figure 1: A representation of $\mathbb{T}_3^F$ with $M_1 = M_2 = 3$ and $M_3 = 2$, with coin tossing as well as waiting time parameters represented beside a few vertices, respectively.

chain of Z_k^F from x is the uniform distribution on the descendants of an ancestor of x whose generation is randomly chosen according to the distribution of g_x.

Remark 2.1. *We may understand the random variable* g_x *as follows. Let us attach coins to the sites of the tree that are not leaves, namely, the points of* $\cup_{j=0}^{k-1}\mathcal{M}|_j = \mathbb{T}_k^F \setminus \mathcal{M}|_k$*, in such a way that the probability of heads of the coin at site* $x|_j \in \mathcal{M}|_j$ *is* $p_j^F(x|_j)$*,* $j = 1, \dots, k-1$*. See Figure 1. The coin of the root has probability* $p_0^F = 0$ *of turning up heads. When it decides to jump from site* $x \in \mathcal{M}|_k$*,* Z_k^F *firstly flips sucessively the coins of* $x|_{k-1}$*, …,* $x|_1$*,* $\emptyset$ *(in that order) until it gets tails for the first time, and then it stops at the respective site. Notice that this procedure is almost surely well defined since* $p_0^F = 0$*. Given that* $x|_j$ *was the stopping site of the procedure, then* $g_x = j$*.*

Definition 2.2. *We call* Z_k^F *a trap model on* $\mathbb{T}_k^F$*, or k-level trap model, with* waiting time *parameter* γ_k^F *and* activation *parameters* $(p_j^F)_{j=1}^{k-1}$*, and write*

$$Z_k^F \sim TM(\mathbb{T}_k^F; \gamma_k^F; (p_j^F)_{j=1}^{k-1}).$$

We are interested in taking limits of Z_k^F as $M_1, \dots, M_k \to \infty$ (under appropriate conditions on its parameters). We will not do this directly but first introduce (in Section 3) a process X_k^F on the leaves of $\mathbb{T}_k^F$, with a certain set of parameters, that coincides with Z_k^F for a specific choice of these parameters.

In Section 4, we define a process X_k, this time on the leaves of the infinite tree $\mathbb{T}_k$, in other words on $\bar{\mathbb{N}}_*^k$, which is an extension of X_k^F to infinite volume. In Section 5, we let $M_i = M_i^{(n)}$, $i = 1, \dots, k$, $n \geq 1$, with $M_i^{(n)} \to \infty$ as $n \to \infty$ for all i. We then consider the process $X_k^{(n)}$ of Section 3, with the "(n)" superscript replacing "F" to emphasize the dependence on n, letting its parameters depend on n as well. Then, under certain conditions on the asymptotics of those parameters as $n \to \infty$, we show that the distribution of $X_k^{(n)}$ converges to that of X_k.

Finally, in Section 6, we consider a particular case of X_k^F, the GREM-like trap model, where the parameters are defined in terms of random variables with heavy tail. Then, by considering the appropriate scaling of the process, including a particular scaling relationship among the volumes $M_1, \dots, M_k$, we establish its convergence to an appropriate X_k by bringing ourselves to the situation of Section 5, and verifying the conditions of the convergence theorem therein.

3 A representation of the k-level trap model

Given $M_1, \dots, M_k \in \mathbb{N}_*$, and recalling the notation of Section 2 above, let $\gamma_j^F : \mathcal{M}|_j \to (0, \infty)$, $j = 1, \dots, k$, and

$$\bar{\gamma}_j^F(x|_j) := \gamma_1^F(x|_1) \times \gamma_2^F(x|_2) \times \dots \times \gamma_j^F(x|_j). \tag{3.1}$$

In this section we will construct a process X_k^F on $\mathcal{M}|_k$ in a inductive way. We first define the process X_1^F. It is a continuous time Markov chain on $\mathcal{M}|_1 (= \mathcal{M}_1)$ that, when at $x_1 \in \mathcal{M}|_1$, waits an exponential time of mean $\gamma_1^F(x_1)$ and then jumps uniformly to a site in $\mathcal{M}|_1$. We will construct X_1^F in the following way.

Let $\mathcal{N}_1 = \{(N_r^{(x_1,1)})_{r \geq 0},\, x_1 \in \mathbb{N}_*\}$ be i.i.d. Poisson processes of rate 1, and let $\sigma_i^{x_1,1}$ be the i-th mark of $N^{(x_1,1)}$ (viewed as a point process), $i \geq 1$. We will call $\mathcal{S}_1^F = \{\sigma_i^{(x_1,1)};\, x_1 \in \mathcal{M}_1,\, i \geq 1\}$ the set of *marks of the first level* of X_k^F. Let $\mathcal{T}_1 = \{T_s^{(1)},\, s \in \mathbb{R}^+\}$ be i.i.d. exponential random variables of rate 1. $\mathcal{N}_1$ and $\mathcal{T}_1$ are assumed independent.

For $s \in \mathcal{S}_1^F$, let $\xi_1^F(s) = x_1$ if $s = \sigma_j^{x_1,1}$ for some $x_1 \in \mathcal{M}_1$ and $j \geq 1$. Notice that ξ_1^F is well defined almost surely. Let us now define a measure μ_1^F on $\mathbb{R}^+ := [0, \infty)$ as follows: $\mu_1^F(\{s\}) = \gamma_1^F(\xi_1^F(s))\, T_s^{(1)}$ if $s \in \mathcal{S}_1^F$, and $\mu_1^F(\mathbb{R}^+ \setminus \mathcal{S}_1^F) = 0$.

Remark 3.1. *We note that $\xi_1^F(s)$, $s \in \mathcal{S}_1^F$, are i.i.d. uniforms in $\mathcal{M}_1$.*

For $r \geq 0$, let

$$\Gamma_1^F(r) := \mu_1^F([0, r]). \tag{3.2}$$

For $t \geq 0$, let

$$\varphi_1^F(t) := (\Gamma_1^F)^{-1}(t) = \inf\{r \geq 0;\, \Gamma_1^F(r) > t\} \tag{3.3}$$

be the (right continuous) inverse of Γ_1^F.

We define the process (X_1^F, Y_1^F) on $(\bar{\mathbb{N}}_*, \mathbb{R}^+)$ as follows. For $t \geq 0$

$$(X_1^F, Y_1^F)(t) = (\xi_1^F(\varphi_1^F(t)), \varphi_1^F(t)). \tag{3.4}$$

Let us suppose (X_j^F, Y_j^F) is defined for $j = 1, \dots, l-1$, $l \leq k$.

Definition 3.2. *An interval $I \subset \mathbb{R}^+$ is a constancy interval of (X_j^F, Y_j^F) if (X_j^F, Y_j^F) is constant over I, that is,*

$$(X_j^F, Y_j^F)(r) = (X_j^F, Y_j^F)(s) \textit{ for all } r, s \in I, \tag{3.5}$$

and I is maximal with that property.



Figure 2: Representation of the timelines of the Poisson point processes entering the definition of X_l^F, one for each $x \in \mathcal{M}_l$. Another ingredient are the constancy intervals of X_{l-1}^F, here successively represented in the x-axis as $I_1, \dots, I_5$.

The maximality condition and right continuity of (X_j^F, Y_j^F) implies that $I = [a, b)$ for some $0 \le a < b$. We are now ready to define (X_l^F, Y_l^F) for $2 \le l \le k$.

Let $\mathcal{I}_{l-1}^F$ be the collection of constancy interval of (X_{l-1}^F, Y_{l-1}^F). Let $\mathcal{N}_l = \{(N_r^{(x_l,l)})_{r\ge0},\ x_l \in \mathbb{N}_*\}$ be i.i.d. Poisson processes of rate 1. Let $\sigma_i^{x_l,l}$ the i-th mark of $N^{(x_l,l)}$, $i \ge 1$. We will call $\mathcal{S}_l^F = \{\sigma_i^{(x_l,l)};\ x_l \in \mathcal{M}_l,\ i \ge 1\}$ the set of Poisson marks of the l-th level, and $\mathcal{R}_l^F = \{a;\ I = [a, b)$ and $I \in \mathcal{I}_{l-1}^F\}$ the set of extra marks of the l-th level. Notice that $\mathcal{R}_l^F$ is the set of left endpoints of intervals of $\mathcal{I}_{l-1}$. We call $\mathcal{S}_l^F \cup \mathcal{R}_l^F$ the set of marks of the l-th level. See Figure 2. Let $\mathcal{T}_l = \{T_s^{(l)},\ s \in \mathbb{R}^+\}$ be i.i.d. exponential random variables of rate 1. $\mathcal{N}_l$ and $\mathcal{T}_l$ are assumed independent and are independent of $\mathcal{N}_j$ and $\mathcal{T}_j$ for $j < l$.

To each $s \in \mathcal{R}_l^F$, we associate an uniform random variable $U_l(s)$ on $\{1, \dots, M_l\}$. Assume that $\{U_l(s),\ s \in \mathcal{R}_l^F,\ l \ge 1\}$ are mutually independent and independent of the other random variables in the model. Let

$$\xi_l^F(s) = \begin{cases} x_l, & \text{if } s = \sigma_j^{(x_l,l)} \text{ for some } x_l \in \mathcal{M}_l \text{ and } j \ge 1, \\ U_l(s), & \text{if } s \in \mathcal{R}_l^F. \end{cases} \tag{3.6}$$

We will call $\xi_l^F(s)$ the *label* of $s \in \mathcal{R}_l^F \cup \mathcal{S}_l^F$, $1 \le l \le k$ (where $\mathcal{R}_1^F = \emptyset$).

Notice that in the first case of (3.6) above $s \in \mathcal{S}_l^F$, and that ξ_l^F is well defined almost surely. Let us now define a measure μ_l^F on $\mathbb{R}^+$ as follows:

$$\mu_l^F(\{s\}) = \gamma_l^F\big(X_{l-1}^F(s), \xi_l^F(s)\big)\, T_s^{(l)}, \ \text{ if } s \in \mathcal{S}_l^F \cup \mathcal{R}_l^F, \tag{3.7}$$



Figure 3: Ilustration of construction of X_k^F from constancy intervals $I_1 = [a_1, b_1), \ldots, I_5 = [a_5, b_5)$ of X_{k-1}^F, on x-axis, Poisson marks, and extra marks, here represented as crosses on left endpoints of the constancy intervals, with (not shown) labels $U_l(a_1), \ldots, U_l(a_5)$, respectively. The Poissonian points may be seen as projected down from timelines of Figure 2. The point "s" in the x-axis equals $\varphi_l^F(t)$.

and $\mu_l^F(\mathbb{R}^+ \setminus (\mathcal{S}_l^F \cup \mathcal{R}_l^F)) = 0$. Notice that $\mathcal{S}_l^F \cap \mathcal{R}_l^F = \emptyset$ almost surely.

Remark 3.3. *We note that* $\xi_j^F(s), s \in \mathcal{S}_j^F \cup \mathcal{R}_j^F$, *are i.i.d. uniforms in* $\mathcal{M}_j, j = 1, \ldots, l$.

For $r \geq 0$, let

$$\Gamma_l^F(r) := \mu_l^F([0, r]). \tag{3.8}$$

For $t \geq 0$, let

$$\varphi_l^F(t) := (\Gamma_l^F)^{-1}(t) = \inf\{r \geq 0 : \Gamma_l^F(r) > t\} \tag{3.9}$$

be the inverse of Γ_l^F.

We define the process (X_l^F, Y_l^F) on $(\mathcal{M}|_l, \mathbb{R}_+^l)$ as follows. For $t \geq 0$

$$(X_l^F, Y_l^F)(t) = \left(\left(X_{l-1}^F(\varphi_l^F(t)), \xi_l^F(\varphi_l^F(t))\right); \left(Y_{l-1}^F(\varphi_l^F(t)), \varphi_l^F(t)\right)\right). \tag{3.10}$$

See Figure 3.

Remark 3.4. *We note that each interval I of $\mathcal{I}_j^F, j \geq 2$, can be identified with a jump of Γ_j^F, that is, $\mathcal{I}_j^F = \{[\Gamma_j^F(r-), \Gamma_j^F(r)) \neq \emptyset;\, r \geq 0\} = \{I_j^F(s) := [\Gamma_j^F(s-), \Gamma_j^F(s));\, s \in \mathcal{S}_j^F \cup \mathcal{R}_j^F\}$, and the lengths of the intervals of $\mathcal{I}_j^F$, namely $\{|I_j^F(s)|,\, s \in \mathcal{S}_j^F \cup \mathcal{R}_j^F\}$, are independent exponential random variables with means $\{\gamma_j^F(X_{j-1}^F(s), \xi_j^F(s)),\, s \in \mathcal{S}_j^F \cup \mathcal{R}_j^F\}$, respectively.*

At this point we may observe that our interest is in X_k^F. Y_k^F is introduced solely for the convenience of having the property mentioned in Remark 3.4. In Section 4 below we will introduce an infinite volume version of X_k^F for which the respective version of Y_k^F will not be needed explicitly, and thus not explicitly introduced.

Remark 3.5. *We note that the number of marks of $\mathcal{S}_j^F$ in each interval $I \in \mathcal{I}_{j-1}^F$ is a geometric random variable with mean $M_j\gamma_{j-1}^F(X_{j-1}^F(s))$, where $s \in \mathcal{S}_{j-1}^F \cup \mathcal{R}_{j-1}^F$ is such that $I = I_{j-1}^F(s)$, and that each such interval has exactly one mark of $\mathcal{R}_j^F$, at its left end. So, the total number of marks (of $\mathcal{S}_j^F \cup \mathcal{R}_j^F$) within I is the above mentioned geometric variable plus one.*

Definition 3.6. *We call X_k^F defined above the trap model on $\mathbb{T}_k^F$, or k-level trap model, with parameter set $\underline{\gamma_k^F} = \{\gamma_i^F;\, i = 1, \ldots, k\}$. Notation: $X_k^F \sim TM(\mathbb{T}_k^F; \underline{\gamma_k^F})$.*

We now make the connection between the models of Section 2 and Section 3, which in particular establishes that the latter is a representation of the former under the appropriate relationship of their respective set of parameters, thus justifying the common terminology.

Lemma 3.7. *Let X_k^F be as above, and Z_k^F be as in Section 2, that is,*

$$Z_k^F \sim TM(\mathcal{M}|_k; \gamma_k^F, (p_j^F)_{j=1}^{k-1}).$$

Suppose

$$p_j^F(x|_j) := \frac{1}{1 + M_{j+1}\gamma_j^F(x|_j)} \tag{3.11}$$

for all $x|_j \in \mathcal{M}|_j$ and $j = 1, \ldots, k-1$. Then X_k^F and Z_k^F have the same distribution.

Proof. We begin with the following remark concerning Z_k^F, which follows immediately from the construction of that process in Section 2.

Remark 3.8. *Let $Z_{k,i}^F$, $i = 1, \ldots, k$, be the i-th coordinate of $Z_k^F = (Z_{k,1}^F, \ldots, Z_{k,k}^F)$, and let $Z_k^F|_j = (Z_{k,1}^F, \ldots, Z_{k,j}^F)$, $j = 1, \ldots, k$. As pointed out in Section 2 above, the jump chain of Z_k^F, let us call it $J_k^F = (J_{k,1}^F, \ldots, J_{k,k}^F)$, with $J_k^F|_j = (J_{k,1}^F, \ldots, J_{k,j}^F)$, $j = 1, \ldots, k$, can be described in terms of the sucessive flips of the coins of $J_k^F|_{k-1}$, …, $J_k^F|_1$ (see Remark 2.1). Let us consider the event $A_{k-1} = \{$flip of the coin of $J_k^F|_{k-1}$ is the first to turn up heads$\}$. In terms of the random variable $g_{J_k^F}$, we have $A_{k-1} = \{g_{J_k^F} = k-1\}$. We now remark that, given A_{k-1}, the distribution of the jump from $J_k^F|_{k-1}$ is the same as that from J_{k-1}^F. Since the distribution of a jump from $J_{k,k}^F$ is always uniform in $\mathcal{M}_k$, we have that, given A_{k-1}, the distribution of the jump from J_k^F is the same as the joint distribution of the jump from J_{k-1}^F, and an independent uniform variable in $\mathcal{M}_k$.*

For $k = 1$, we remark that the transition probabilities of X_1^F are always uniform in $\mathcal{M}_1$, since the labels of the sucessive points of $\mathcal{S}_1^F$ have this property and are independent of each other. So, X_1^F and Z_1^F are processes with the same state space and transition probabilities, and the holding times are clearly matched. The result follows for $k = 1$.

We now proceed by induction on k. Let us suppose that the result holds for $k = K - 1$ for some $K \geq 2$. To show that $X_K^F \sim Z_K^F$, it is enough to identify the transition mechanisms of both processes. Let us thus fix a time $t \geq 0$ and a point $x|_K \in \mathcal{M}|_K$. Given that either process is at $x|_K$ at time t, then both jump times are exponentially distributed with mean $\gamma_K^F(x|_K)$. So far, we have an identification. Now let us identify the jump mechanisms of both processes.

As discussed in Remark 3.8 above, given A_{K-1} we have that $Z_K^F|_{K-1}$ jumps as Z_{K-1}^F, and $Z_{K,K}^F$ jumps uniformly in $\mathcal{M}_K$, and the jumps of $Z_K^F|_{K-1}$ and $Z_{K,K}^F$ are independent. Let us identify a *coin tossing* mechanism in the jump of X_K^F. Let I be the interval of $\mathcal{I}_{K-1}^F$ containing $\varphi_K(t)$, and let $I' = (\varphi_K(t), \infty) \cap I$. The lack of memory of the exponential distribution implies that, given that $X_K^F(t) = x|_K$, then $|I'|$ is an exponential random variable with mean $\gamma_{K-1}^F(x|_{K-1})$. The mechanism in the X_K^F process that plays the role of (the first) coin tossing is whether or not I' contains at least one (Poisson) mark. Let us call $\bar{A}_{K-1}$ the event that I' contains no mark. This corresponds to the coin turning up heads. This means that $X_K^F|_{K-1}$ will take a jump, which we identify as a jump of X_{K-1}^F, independent of the (uniform in $\mathcal{M}_K$) accompanying jump of $X_{K,K}^F$.

At this point we should stress that a particular element of the construction of X_K^F plays a key role in this identification, namely, the inclusion of the marks of $\mathcal{R}_K^F$. This guarantees that each interval of constancy I of $\mathcal{I}_{k-1}^F$ gets at least one mark. Without these marks, the jump of $X_K^F|_{K-1}$ might not coincide with that of X_{K-1}^F – that would happen if (and only if) the first interval of $\mathcal{I}_{K-1}^F$ neighboring I to the right had got no mark.

By the induction hypothesis, $X_{K-1}^F \sim Z_{K-1}^F \sim Z_K^F|_{K-1}$; clearly $X_{K,K}^F \sim Z_{K,K}^F$. We close the argument in two steps. The first one is to show that

$$P\big(\bar{A}_{K-1}\,\big|X_K^F(t) = x|_K\big) = \frac{1}{1 + M_K \gamma_{K-1}^F(x|_{K-1})} = P\big(A_{K-1}\,\big|Z_K^F(t) = x|_K\big)\,, \tag{3.12}$$

and then use the hypothesis. But it follows from our discussion above and the construction of X_K^F that the left hand side of (3.12) equals $P(N'_{T'} = 0)$, where N' is a Poisson process of rate M_k and T' an independent exponential random variable with mean $\gamma_{K-1}^F(x|_{K-1})$, and a simple computation yields the result.

And the last step is to argue that, given $\bar{A}_{K-1}^c$, then $X_K^F|_{K-1}$ does not move, and we have only the uniform in $\mathcal{M}_K$ jump of $X_{K,K}^F$, which agrees with the corresponding move of Z_K^F given A_{K-1}^c.

□

4 K process on a tree

K processes on $\bar{\mathbb{N}}_*$ were introduced in [9] in the study of limits of trap models in the complete graph. They appear as scaling limits of the REM-like trap model in the complete graph. Below we introduce an extension of that model to a model on $\bar{\mathbb{N}}_*^k$, which we will view as the leaves of a tree with k generations, similarly as done in the previous sections.The aim is, similarly as in [9], to establish limit results for the processes of the previous sections as volume diverges.

Let $\gamma_j : \mathbb{N}_*^j \to (0, \infty)$, $j = 1, \ldots, k$, be such that, making

$$\bar{\gamma}_j(x|_j) := \gamma_1(x|_1) \times \gamma_2(x|_2) \times \ldots \times \gamma_j(x|_j), \tag{4.1}$$

we have

$$\sum_{x|_j \in \mathbb{N}_*^j} \bar{\gamma}_j(x|_j) < \infty \tag{4.2}$$

We will construct a process X_k on $\bar{\mathbb{N}}_*^k$ inductively, similarly as in Section 3. First we define the process X_1. It is a continuous time Markov chain on $\bar{\mathbb{N}}_*$ described as follows.

Let $\mathcal{N}_1 = \{(N_r^{(x_1,1)})_{r \geq 0},\, x_1 \in \mathbb{N}_*\}$ be i.i.d. Poisson processes of rate 1. Let $\sigma_i^{x_1,1}$ be the i-th mark of $N^{(x_1,1)}$, $i \geq 1$. We will call $\mathcal{S}_1 = \{\sigma_i^{(x_1,1)};\, x_1 \in \mathbb{N}_*,\, i \geq 1\}$ the set of marks of the first level of X_k. Let $\mathcal{T}_1 = \{T_s^{(1)},\, s \in \mathbb{R}^+\}$ be i.i.d. exponential random variables of rate 1. $\mathcal{N}_1$ and $\mathcal{T}_1$ are assumed independent.

For $s \in \mathcal{S}_1$, let $\xi_1(s) = x_1$ if $s = \sigma_i^{x_1,1}$ for some $x_1 \in \mathbb{N}_*$ and $i \geq 1$. Notice that ξ_1 is well defined almost surely. Let us now define a measure μ_1 on $\mathbb{R}^+$ as follows.

$$\mu_1(\{s\}) = \gamma_1(\xi_1(s))\, T_s^{(1)}, \text{ if } s \in \mathcal{S}_1, \text{ and } \mu_1(\mathbb{R}^+ \setminus \mathcal{S}_1) = 0. \tag{4.3}$$

For $r \geq 0$, let

$$\Gamma_1(r) := \mu_1([0, r]), \tag{4.4}$$

and, for $t \geq 0$, let

$$\varphi_1(t) := \Gamma_1^{-1}(t) = \inf\{r \geq 0 : \Gamma_1(r) > t\} \tag{4.5}$$

be the inverse of Γ_1.

Remark 4.1. *Notice that μ_1 is almost surely a purely atomic measure whose set of atoms, $\mathcal{S}_1$, is a.s. countable and dense in $\mathbb{R}^+$. Moreover, from Lemma 4.5 below, it is a.s. σ-finite. These properties imply that $\Gamma_1 : \mathbb{R}^+ \to \mathbb{R}^+$ is a.s. strictly increasing and that its range $\Gamma_1(\mathbb{R}^+)$ is an uncountable set of Lebesgue measure zero. It follows from this and the independence and continuity of its constituents that any fixed deterministic r is a.s. a continuity point of Γ_1. It may also be checked that $\varphi_1 : \mathbb{R}^+ \to \mathbb{R}^+$ is a.s. continuous.*

We define X_1 on $\bar{\mathbb{N}}_*$ as follows. For $t \geq 0$

$$X_1(t) = \begin{cases} \xi_1(\varphi_1(t)), \text{ if } \varphi_1(t) \in \mathcal{S}_1, \\ \infty, \text{ otherwise.} \end{cases} \tag{4.6}$$

Let us suppose X_j is defined for $j = 1, \ldots, l-1$, $2 \leq l \leq k$.

Definition 4.2. *An interval $I \subset \mathbb{R}^+$ is a constancy interval of X_j if it has positive length,*

$$X_j(r) = X_j(s) \textit{ for all } r, s \in I, \textit{ and } I \textit{ is maximal.} \tag{4.7}$$

The maximality condition and right continuity of X_j implies that $I = [a, b)$ for some $0 \leq a < b$. We are now ready to define X_l.

Let $\mathcal{I}_{l-1}$ be the collection of constancy interval of X_{l-1}. Let $\mathcal{N}_l = \{(N_r^{(x_l,l)})_{r\geq 0},\, x_l \in \mathbb{N}_*\}$ be i.i.d. Poisson processes of rate 1. Let $\sigma_i^{x_l,l}$ the i-th mark of $N^{(x_l,l)}$, $i \geq 1$. We will call $\mathcal{S}_l = \{\sigma_i^{(x_l,l)};\, x_l \in \mathbb{N}_*,\, i \geq 1\}$ the set of Poisson marks of the l-th level. Let $\mathcal{T}_l = \{T_s^{(l)},\, s \in \mathbb{R}^+\}$ be i.i.d. exponential random variables of rate 1. $\mathcal{N}_l$ and $\mathcal{T}_l$ are assumed independent and are independent of $\mathcal{N}_j$ and $\mathcal{T}_j$ for $j < l$.

For $s \in \mathcal{S}_l$, let $\xi_l(s) = x_l$ if $s = \sigma_j^{(x_l,l)}$ for some $x_l \in \mathbb{N}_*$ and $j \geq 1$. Notice that ξ_l is well defined almost surely. Let us now define a measure μ_l on $\mathbb{R}^+$ as follows:

$$\mu_l(\{s\}) = \gamma_l(X_{l-1}(s), \xi_l(s))T_s^{(l)}, \text{ if } s \in \mathcal{S}_l, \text{ and } \mu_l(\mathbb{R}^+ \setminus \mathcal{S}_l) = 0.$$

For $r \geq 0$, let

$$\Gamma_l(r) := \mu_l([0, r]), \tag{4.8}$$

and for $t \geq 0$, let

$$\varphi_l(t) := \Gamma_l^{-1}(t) = \inf\{r \geq 0;\, \Gamma_l(r) > t\} \tag{4.9}$$

be the inverse of Γ_l.

Remark 4.3. *Remark 4.1 holds with "1" replaced by "l". In particular, the range of Γ_l has a.s. Lebesgue measure zero, $l = 1, \ldots, k$, and every fixed deterministic r is a.s. a continuity point of Γ_l.*

We define the process X_l on $\bar{\mathbb{N}}_*^l$ as follows. For $t \geq 0$, let

$$X_l(t) = \begin{cases} (X_{l-1}(\varphi_l(t)), \xi_l(\varphi_l(t)))\,, \text{ if } \varphi_l(t) \in \mathcal{S}_l, \\ (X_{l-1}(\varphi_l(t)), \infty)\,, \text{ otherwise.} \end{cases}$$

Definition 4.4. *We call X_k defined just above the K process on $\mathbb{T}_k$, or k-level K process, with parameter set $\underline{\gamma_k} = \{\gamma_i;\, i = 1, \ldots, k\}$. Notation: $X_k \sim K(\mathbb{T}_k, \underline{\gamma_k})$.*

Pictures like those in Figures 2 and 3 might be drawn (or perhaps, more accurately, envisioned) for X_k, with minor changes: in the present case we would have an infinite sequence of timelines for the Poisson processes $(N_r^{(x_l,l)})_{r\geq 0}$, $x_l \in \mathbb{N}_*$, in Figure 2. In Figure 3, superscripts "F" should be dropped throughout; there would be no extra marks, and thus no crosses; the Poissonian marks would form a dense set of the x-axis; the constancy intervals should be such that there would be an infinite number of them in the neighborhood of any fixed one of them – or, more precisely, between any two distinct such intervals, there is a distinct such interval; the graph of Γ_l would be that of a strictly increasing function with a dense set of jumps (the Poissonian marks).

The next result makes the above construction well-defined. For its proof, let us introduce the notation

$$X_k = (X_{k,1}, \ldots, X_{k,k}),\ k \geq 1,$$

making the coordinates of X_k explicit.

Lemma 4.5. *Almost surely* $\Gamma_k(r) < \infty$ *for all* $r \in (0, \infty)$.

Proof. As Γ_j is nondecreasing and unbounded for $j = 1, \ldots, k$, it is suficient to show that, for all $r \in (0, \infty)$, $\Gamma_k \circ \ldots \circ \Gamma_1(r) < \infty$ almost surely. Let

$$\Theta_k(r) := \Gamma_k \circ \ldots \circ \Gamma_1(r). \tag{4.10}$$

We will show by induction that

$$E(\Theta_k(r)) = r \sum_{x_1=1}^{\infty} \ldots \sum_{x_k=1}^{\infty} \gamma_1(x|_1) \ldots \gamma_k(x|_k) = \sum_{x \in \mathbb{N}_*^k} \bar{\gamma}_k(x), \tag{4.11}$$

and this is finite by assumption (see (4.2) above), thus closing the argument.

This is immediate from the definition for $k = 1$. Let us suppose that it holds up to $k - 1$, $k \geq 2$. Let us consider the constancy intervals of $X_{k,1}$ (that is, maximal intervals over which $X_{k,1}$ is constant): $\mathcal{I} = \{$constancy intervals of $X_{k,1} \subset [0, \Theta_k(r)]\}$. We can enumerate such intervals as $\mathcal{I} = \{I(s);\, s \in \mathcal{S}_1 \cap [0, r]\}$. So,

$$\Theta_k(r) = \sum_{s \in \mathcal{S}_1 \cap [0,r]} |I(s)| = \sum_{x_1=1}^{\infty} \sum_{i_1=1}^{N_r^{(x_1)}} L_{i_1}^{(x_1)}, \tag{4.12}$$

where $L_{i_1}^{(x_1)} := |I(\sigma_{i_1}^{(x_1,1)})|$. Now notice that, for every $x_1 \in \mathbb{N}_*$, $N^{(x_1)}$ and $L^{(x_1)} := \{L_{i_1}^{(x_1)} : i_1 \geq 1\}$ are independent, and $L^{(x_1)}$ is an i.i.d. family of random variables with $L_{i_1}^{(x_1)} \sim \Theta_{k-1}^{(x_1)}(\gamma_1(x_1) T_1^{(x_1)})$, where $\Theta_{k-1}^{(x_1)} = \Gamma_{k-1} \circ \ldots \circ \Gamma_1$ is the corresponding of Θ_k for a K process on $\mathbb{T}_k^{(x_1)}$, the $k-1$-level subtree of $\mathbb{T}_k$ rooted on x_1, and parameter set $\underline{\gamma_k^{(x_1)}} = \{\gamma_i(x_1, \cdot) : i = 2, \ldots, k\}$.

Then,

$$E(\Theta_k(r)) = r \sum_{x_1=1}^{\infty} E\{\Theta_{k-1}^{(x|_1)}(\gamma_1(x|_1)\, T_1^{(x_1)})\} = r \sum_{x_1=1}^{\infty} \gamma_1(x|_1) \sum_{x_2, \ldots, x_k} \gamma_2(x|_2) \ldots \gamma_k(x|_k), \tag{4.13}$$

where we have used that $T_{i_1}^{(x_1)}$, $i_1 \geq 1$, are i.i.d. with mean 1 random variables, independent of all other random variables, and, in the second equality, the induction hypothesis. The coincidence of the right hand sides of (4.11) and (4.13) closes the argument. □

Remark 4.6. *It follows readily from (4.12) and the independence of the summands on its right hand side, and the fact that they are identically distributed, that a.s.* $\Theta_k(r) \to \infty$ *as* $r \to \infty$.

Remark 4.7.

1. *We further notice that* $X_{j,j}(t) = \infty$ *for a given* t *if and only if* $t \in \Gamma_j(\mathbb{R}_+)$.

2. *From Remarks 4.3 and the previous item, it follows that for* $j = 1, \ldots, k$, *the infinities of* X_j, *that is, the set* $\{t \geq 0 : X_{j,i}(t) = \infty$ *for some* $i = 1, \ldots, j\}$ *has a.s. Lebesgue measure zero.*

3. It follows that a.s. $X_{j,i}(s) < \infty$ *for* $s \in \mathcal{S}_{j+1}$, $i = 1, \ldots, j$, *and* $j = 1, \ldots, k-1$.

4. It follows that a.s. if $X_{k,i}(t) = \infty$ *for some* $t \geq 0$ *and* $i = 1, \ldots, k$, *then* $X_{k,j}(t) = \infty$ *for* $i \leq j \leq k$.

The next result states roughly that once a coordinate of a K process is large, then so are the subsequent ones. This is in line with the property mentioned in Remark 4.7.4 above – a sort of continuous extension of it. In the next section we establish a finite volume analogue (see Lemma 5.7 below).

Lemma 4.8. *Let* X_k *be a* k*-level K process. Given* $T > 0$ *and* $m \geq 1$, *there a.s. exist* $\tilde{m} = \tilde{m}^{(k)}$ *such that if* $X_{k,i}(t) > \tilde{m}$ *and* $t \in [0, T]$, *then* $X_{k,j}(t) > m$ *for all* $j = i+1, \ldots, k$.

Proof. Let $T' > 0$ be such that $\Theta_k(T') > T$ (see Remark 4.6), and for $m \in \mathbb{N}_*$ and $j = 1. \ldots, k$, let

$$\tilde{\mathcal{S}}_j^{(m)} = \{\sigma_i^{(x,j)} : x = 1, \ldots, m, \, i \geq 1\} \tag{4.14}$$

be the set of Poissonian marks of level j with labels at most m. Let $\mathcal{T}_{kj}(l)$ denote the set of times up to $\Theta_{k-1}(T')$ spent by $X_{k-1,j}$ above l, $j = 1, \ldots, k-1$. By a similar reasoning as the one employed to prove (4.11), we may check that the expected length of $\mathcal{T}_{kj}(l)$ equals

$$T' \sum_{x_1=1}^{\infty} \ldots \sum_{x_j=l+1}^{\infty} \ldots \sum_{x_{k-1}=1}^{\infty} \gamma_1(x|_1) \ldots \gamma_{k-1}(x|_{k-1}), \tag{4.15}$$

and since that length is decreasing in l, and, as follows from our assumptions on $\underline{\gamma_k}$, the expression in (4.15) vanishes as $l \to \infty$, we have that the limit of that length as $l \to \infty$ vanishes almost surely. We then have from elementary properties of Poisson processes that

$$\{\cup_{j=1}^{k-1} \mathcal{T}_{kj}(l)\} \cap \tilde{\mathcal{S}}_k^{(m)} = \emptyset \tag{4.16}$$

for all large enough l almost surely, so given $m \in \mathbb{N}_*$, we find $\hat{m}^{(k)}$ such that if $X_{k,i}(t) > \hat{m}^{(k)}$ for any $t \leq T$ and $i = 1, \ldots, k-1$, then it follows that $X_{k,k}(t) > m$. This in particular establishes the claim for $k = 2$ by the choice $\tilde{m}^{(2)} = \hat{m}^{(2)}$. Let us assume that the claim is established for $k-1$. Then substituting $\Theta_{k-1}(T')$ as T, and $\hat{m}^{(k)} \vee m$ as m in that claim, and taking the choice $\tilde{m}^{(k)} = \tilde{m}^{(k-1)} \vee \hat{m}^{(k)}$, we find that it satisfies the claim.

□

5 Convergence

5.1 Statement of results

In this section, we consider a sequence of trap models $X_k^{(n)}$, $n \geq 1$, on a sequence of finite trees $\mathbb{T}_k^{(n)}$, with volumes $M_1 = M_1^{(n)}, \ldots, M_k = M_k^{(n)}$, and parameter sets $\underline{\gamma_k^{(n)}}$, respectively, and prove a weak convergence result for that sequence under the Skorohod topology on $D(\bar{\mathbb{N}}_*^k, [0, \infty))$, the space of càdlàg functions from $[0, \infty)$ to $\bar{\mathbb{N}}_*^k$. As anticipated at the end of Section 3, we will replace the superscript "F" by "(n)" everywhere to emphasize the dependence on n.

Before proceeding, let us briefly review this topology. We start by equipping $\bar{\mathbb{N}}_*^k$ with the metric

$$d(x,y) = \max_{1\leq j\leq k} |x_j^{-1} - y_j^{-1}|,\ x,\, y \in \bar{\mathbb{N}}_*^k, \tag{5.1}$$

where $\infty^{-1} = 0$, under which it is compact. The Skorohod metric on $D(\bar{\mathbb{N}}_*^k, [0,\infty))$ is as follows. For $f, g \in D(\bar{\mathbb{N}}_*^k, [0,\infty))$, let

$$\rho(f,g) = \inf_{\lambda\in\Lambda} \left[\phi(\lambda) \vee \int_0^\infty e^{-u} \rho(f,g,\lambda,u)\, du\right], \tag{5.2}$$

where

$$\rho(f,g,\lambda,u) = \sup_{t\geq 0} d(f(t\wedge u), g(\lambda(t)\wedge u)), \tag{5.3}$$

with Λ the class of *time distortions*: increasing Lipschitz functions from $[0,\infty)$ onto $[0,\infty)$, and $\phi : \Lambda \to [0,\infty)$ such that

$$\phi(\lambda) = \sup_{0\leq s<t} \left|\log \frac{\lambda_t - \lambda_s}{t-s}\right| \tag{5.4}$$

(see Section 3.5 in [24]).

In order to get our convergence result, we will impose the following conditions on the volumes and parameters. For $j = 1, \dots, k$, suppose that as $n \to \infty$

$$M_j^{(n)} \to \infty, \tag{5.5}$$

$$\gamma_j^{(n)}(x) \to \gamma_j(x) \text{ for every } x \in \mathbb{N}_*^j, \text{ and } \sum_{x\in\mathbb{N}_*^j} \bar{\gamma}_j^{(n)}(x) \to \sum_{x\in\mathbb{N}_*^j} \bar{\gamma}_j(x), \tag{5.6}$$

with γ_j, $\bar{\gamma}_j$ as in the beginning of Section 4 (see paragraph of (4.1,4.2) above), and $\gamma_j^{(n)} \equiv 0$ on $\mathbb{N}_*^j \setminus \mathcal{M}|_j$.

Our result will require additional conditions, that look quite intricate. We state them now and discuss them, together with the above ones, after we state the convergence result. We further suppose that for $j = 2, \dots, k$

$$\frac{1}{\prod_{p=2}^j M_p} \sum_{l=1}^{j-1} \sum_{x|_{j-1}\in\mathcal{M}|_{j-1}} \prod_{p=1}^{l-1} M_{p+1}\, \gamma_p^{(n)}(x|_p) \prod_{p=l+1}^{j-1} (1 + M_{p+1}\, \gamma_p^{(n)}(x|_p)) \to 0 \tag{5.7}$$

and

$$\frac{1}{\prod_{p=2}^j M_p} \sum_{l=1}^{j-1} \sum_{x|_j\in\mathcal{M}|_j} \gamma_j^{(n)}(x|_j) \prod_{p=1}^{l-1} M_{p+1}\, \gamma_p^{(n)}(x|_p) \prod_{p=l+1}^{j-1} (1 + M_{p+1}\, \gamma_p^{(n)}(x|_p)) \to 0 \tag{5.8}$$

as $n \to \infty$, where by convention

$$\prod_{p=2}^{1} M_q = \prod_{p=1}^{0} M_{p+1}\, \gamma_p^{(n)}(x|_p) = \prod_{p=j}^{j-1} (1 + M_{p+1}\, \gamma_p^{(n)}(x|_p)) = 1.$$

Here, and many times below, we omit the superscript "(n)" from the notation for the volumes $M_1, \dots, M_k$.

We are ready to sate our main result.

Theorem 5.1. *For $n \geq 1$, let $X_k^{(n)}$ be the trap model on $\mathbb{T}_k^{(n)}$, with volumes $M_1^{(n)}, \dots, M_k^{(n)}$, and parameter sets $\underline{\gamma_k^{(n)}}$, respectively, satisfying conditions (5.5-5.8). Let X_k be the K process on $\mathbb{T}_k$ with parameter set $\underline{\gamma_k}$. Then $X_k^{(n)} \to X_k$ weakly in Skorohod space as $n \to \infty$.*

We will see (from the proofs) that conditions (5.5-5.8) have the following significance. Obviously, (5.5) means that we are taking an infinite volume limit. Equation 5.6 implies that the contributions coming from the Poisson marks to the construction of $X_k^{(n)}$ converge (in a uniform way) to the respective contributions of (Poisson) marks of X_k. Finally, as will be seen in the arguments below, (5.7-5.8) imply the negligibility of the total contribution of the extra marks entering $X_k^{(n)}$.

The remainder of this section is organized as follows. We briefly start below, in this same subsection, with the proof of Theorem 5.1. The full proof will require a number of auxiliary results, which we collect in Subsection 5.2 below, before proceeding with the proof in Subsection 5.3 after that.

Proof of Theorem 5.1

We will argue by induction, using coupled versions of $X_k^{(n)}$ and X_k, and show convergence in probability for a subsequence. The coupling is going to be given by using common Poisson processes $\{N^{(x_i,i)}, x_i \in \mathbb{N}_*, i = 1, \dots, k\}$ and common exponential variables $\{T_s^{(i)}, s \in \mathbb{R}^+, i = 1, \dots, k\}$ in the construction of $X_k^{(n)}$ and X_k. The notation is detailed at the beginning of Subsection 5.3. It will be clear that the same can be done for every subsequence of (n), and that the limiting distribution for each subsubsequence does not depend on the subsequence. This then implies weak convergence of the original sequence.

Lemma 3.11 of [9] establishes the (convergence in probability; actually a.s. convergence) result for $k = 1$ and $\gamma_1^{(n)}(x)$ not depending on n as soon as $x \leq M_1$. This result (convergence in probability) holds (with minor changes in argumentation, as sketched in the proof of Theorem 5.2 of [9]) in our case as well. It is also part of the argumentation of Lemma 3.11 and Theorem 5.2 of [9]), and can also be readily checked independently, that for every $r \in [0, \infty)$

$$\Gamma_1^{(n)}(r) \to \Gamma_1(r) \tag{5.9}$$

in probability as $n \to \infty$.

As part of our induction argument, we will the assume that for $j = 1, \dots, k-1$ and every $r \in [0, \infty)$

$$\begin{aligned} X_j^{(n)} &\to X_j && (5.10)\\ \Gamma_j^{(n)}(r) &\to \Gamma_j(r) && (5.11) \end{aligned}$$

as $n \to \infty$ almost surely, possibly over a subsequence.

5.2 Preliminaries

Our first auxiliary result establishes that the contribution of extra marks and their descendants to $X_j^{(n)}$ is negligible as $n \to \infty$. That is the content of Lemma 5.2. To be precise, let $\mathcal{E}_2^{(n)} = \mathcal{R}_2^{(n)}$ and for $3 \leq i \leq k$

$$\mathcal{E}_i^{(n)} = \mathcal{R}_i^{(n)} \cup \{s \in \mathcal{S}_i^{(n)} : \varphi_{i-1}^{(n)}(s) \in \mathcal{E}_{i-1}^{(n)}\}. \tag{5.12}$$

$\mathcal{E}_i^{(n)}$ represents the extra marks of the i-th level and the *descendants* of extra marks of previous levels in the i-th level (that is, Poisson marks belonging to constancy intervals originating from extra marks of the previous level or descendants of extra marks from levels before that).

Lemma 5.2. *Assume that the induction hypotheses (5.10-5.11) hold. Then, for every $r > 0$ and $j = 2, \ldots, k$, we have that*

a) $\min\{\xi_j^{(n)}(s) : s \in \mathcal{E}_j^{(n)} \cap [0, r]\} \to \infty$ *in probability as $n \to \infty$.*

b) $\mu_j^{(n)}(\mathcal{E}_j^{(n)} \cap [0, r]) \to 0$ *in probability as $n \to \infty$.*

Proof. For $r > 0$, $j = 1, \ldots, k$, let

$$\Theta_j^{(n)}(r) := \Gamma_j^{(n)} \circ \ldots \circ \Gamma_1^{(n)}(r), \tag{5.13}$$

and define $K_j^{(n)}(r) := |\mathcal{E}_j^{(n)} \cap [0, \Theta_{j-1}^{(n)}(r)]|$, where (here) $|\cdot|$ stands for cardinality. An evaluation of $K_j^{(n)}(r)$ will play a crucial role in the proof. We begin with that.

It follows from induction hypothesis (5.11) that for $j = 1, \ldots, k-1$, $\Gamma_j^{(n)}(r) \to \infty$ as $r \to \infty$ in probability, uniformly in n. So it is enough to consider $\mathcal{E}_j^{(n)} \cap [0, \Theta_{j-1}^{(n)}(r)]$. In order to evaluate the cardinality of that set, as well as its contribution to $\mu_j^{(n)}([0, \Theta_{j-1}^{(n)}(r)])$, we start by describing the structure of $\mathcal{H}_1^{(n)} := \mathcal{S}_1^{(n)}$, $\mathcal{H}_2^{(n)} := \mathcal{S}_2^{(n)} \cup \mathcal{R}_2^{(n)}$, …, $\mathcal{H}_k^{(n)} := \mathcal{S}_k^{(n)} \cup \mathcal{R}_k^{(n)}$; at the same time, we will relabel the marks of those sets conveniently.

Each $s_1 \in \mathcal{S}_1^{(n)}$ can be put in a one to one correspondence with its label $\xi_1(s_1) = x_1$ and *index* $i_1(s_1) = i_1 \in \mathbb{N}_*$ via the relation $s_1 = \sigma_{i_1}^{(x_1,1)}$. Using this correspondence, we see that to each mark s_1 of $\mathcal{S}_1^{(n)}$ there corresponds an interval $I_n^{(x_1,i_1)}$ of $\mathbb{R}_+$ of length $L_n^{(x_1,i_1)} = \gamma_1^{(n)}(x_1)\, T^{(x_1,i_1)}$. Such intervals form a partition of $\mathbb{R}_+$, and the random variables involved are independent when we vary s_1.

Now to each $s_1 \in \mathcal{S}_1^{(n)}$, there corresponds marks of $\mathcal{S}_2^{(n)}$ belonging to the respective interval $I_n^{(x_1,i_1)}$, whose cardinality is a geometric random variable $G^{(x_1,i_1)}$ with mean $M_2\, \gamma_1^{(n)}(x_1)$, plus a mark of $\mathcal{R}_2^{(n)}$ at the left endpoint of $I_n^{(x_1,i_1)}$. Each such mark will be identified with $(x_1, i|_2)$, where (x_1, i_1) is the identifier of s_1, and $i_2 \in \{1, \ldots, G^{(x_1,i_1)} + 1\}$, and we attach to it a random variable $U^{(x_1,i|_2)}$ with uniform distribution in $\mathcal{M}_2$, which corresponds to $\xi_2(s_2)$, for $(x_1, i|_2) \equiv s_2 \in \mathcal{S}_2^{(n)} \cup \mathcal{R}_2^{(n)}$. We will identify the unique mark of $\mathcal{R}_2^{(n)}$ at the left endpoint of $I_n^{(x_1,i_1)}$ with $(x_1, i_1, 1)$.

We now proceed inductively. For $3 \le j \le k$, we assume we have identified each mark of $\mathcal{S}_{j-1}^{(n)} \cup \mathcal{R}_{j-1}^{(n)}$ as $(x_1, i|_{j-1})$, with $x_1 \in \mathcal{M}_1$, $i_1 \ge 1$, $i_l = 1, \ldots, 1 + G^{(x_1,i|_{l-1})}$, $l = 2, \ldots, j-1$, where, for $l \ge 3$, $G^{(x_1,i|_{l-1})}$ is geometric with mean $M_l\, \gamma_{l-1}^{(n)}(x_1, U^{(x_1,i|_2)}, \ldots, U^{(x_1,i|_{l-1})})$, with $U^{(x_1,i|_j)} \sim \text{Uniform}(\mathcal{M}_j)$. The random variables of

$$\mathcal{U}_{j-1} := \{U^{(x_1,i|_l)} : l = 2, \ldots, j-1;\, x_1, i_1, \ldots, i_l \ge 1\}$$

are independent, and, given $\mathcal{U}_{j-1}$, so are the ones of

$$\{G^{(x_1,i|_l)} : l = 1, \ldots, j-1;\, x_1, i_1, \ldots, i_l \ge 1\}$$

— notice that $G^{(x_1,i|_j)}$ is independent of $U^{(x_1,i|_l)}$ as soon as $j < l$. Here $i_{j-1} = 1$ means $(x_1, i|_{j-1}) \in \mathcal{R}^{(n)}_{j-1}$; otherwise, $(x_1, i|_{j-1}) \in \mathcal{S}^{(n)}_{j-1}$. Then to each mark $(x_1, i|_{j-1})$ there corresponds an interval $I_n^{(x_1,i|_{j-1})}$ of $\mathbb{R}_+$ of length

$$L_n^{(x_1,i|_{j-1})} := \gamma^{(n)}_{j-1}(x_1, U^{(x_1,i|_2)}, \ldots, U^{(x_1,i|_{j-1})})\, T^{(x_1,i|_{j-1})},$$

with $\{T^{(x_1,i|_{j-1})}\}$ i.i.d. mean 1 exponential random variables independent of $\{G^{(x_1,i|_l)}, U^{(x_1,i|_l)}\}$, $l = 1, \ldots, j-1$, such that $\{I_n^{(x_1,i|_{j-1})}\}$ is a partition of $\mathbb{R}_+$. The mark of $\mathcal{R}^{(n)}_j$ placed at the left end of $I_n^{(x_1,i|_{j-1})}$ is labeled $(x_1, i|_{j-1}, 1)$, and the marks of $\mathcal{S}^{(n)}_j \in I_n^{(x_1,i|_{j-1})}$, if any, are labeled $(x_1, i|_j)$, $i_j = 2, \ldots, 1+G^{(x_1,i|_{j-1})}$, with $G^{(x_1,i|_{j-1})}$ a geometric random variable with mean $M_j\, \gamma^{(n)}_{j-1}(x_1, U^{(x_1,i|_2)}, \ldots, U^{(x_1,i|_{j-1})})$. The random variables in $\{G^{(x_1,i|_{j-1})}\}$ are independent and independent of the previous random variables. Finally, we assign to each $(x_1, i|_j)$ a random variable $U^{(x_1,i|_j)}$ uniformly distributed in $\mathcal{M}_j$, corresponding to $\xi_j(s_j)$, for $(x_1, i|_j) \equiv s_j \in \mathcal{S}^{(n)}_j \cup \mathcal{R}^{(n)}_j$, with $\{U^{(x_1,i|_j)}\}$ independent and independent of previous random variables.

With this representation, we have labeled the marks of $\mathcal{H}^{(n)}_j \cap [0, \Theta^{(n)}_{j-1}(r)]$, $j = 1, \ldots, k$, as $(x_1, i|_j)$, $x_1 = 1, \ldots, M_1$; $i_1 = 1, \ldots, N_r^{(x_1)}$; $i_l = 1, \ldots,\ 1 + G^{(x_1,i|_{l-1})}$, $2 \le l \le j$, with $G^{(x_1,i|_l)}$ geometric with mean $M_2\, \gamma^{(n)}_1(x_1)$, when $l = 1$, and $M_{l+1}\, \gamma^{(n)}_l(x_1, U^{(x_1,i_1)}, \ldots, U^{(x_1,i|_l)})$, when $l = 2, \ldots, j$, respectively. $U^{(x_1,i|_l)}$ is uniformly distributed on $\mathcal{M}_l$, $l = 2, \ldots, j$. The random variables in the family

$$\mathcal{U}_j := \{U^{(x_1,i|_l)};\ x_1, i_1, \ldots, i_l \ge 1,\ l = 2, \ldots, j\}$$

are independent, and given $\mathcal{U}_j$ so are the ones in

$$\{G^{(x_1,i|_l)};\ x_1, i_1, \ldots, i_l \ge 1,\ l = 1, \ldots, j\}$$

— notice that, as before, $G^{(x_1,i|_j)}$ is independent of $U^{(x_1,i|_l)}$ as soon as $j < l$.

The marks of $\mathcal{E}^{(n)}_j \cap [0, \Theta^{(n)}_{j-1}(r)]$ are those $(x_1, i|_j)$ as above for which $i_l = 1$ for some $l = 2, \ldots, j$. In order to write an expression for $K^{(n)}_j(r)$ we first view $\mathcal{E}^{(n)}_j \cap [0, \Theta^{(n)}_{j-1}(r)]$ as the leaves of a forest (see Figure 4), the distinct trees of which have the marks labeled $(x_1, i|_l, 1)$, $l = 1, \ldots, j-1$, $i_1 = 1, \ldots, N_r^{(x_1)}$; $i_l = 1, \ldots, \tilde{G}^{(x_1,i|_{l-1})} := 1 + G^{(x_1,i|_{l-1})}$, $l = 2, \ldots, j-1$, as roots; the tree rooted at $(x_1, i|_l, 1)$ consisting of, besides the root, marks whose labels form the set

$$\mathfrak{T}_j^{(x_1,i|_l)} := \{(x_1, i|_l, 1, i_{l+2}, \ldots, i_j),\ i_m = 1, \ldots, \tilde{G}^{(x_1,i|_{m-1})},\ m = l+2, \ldots, j\}. \tag{5.14}$$

The latter set is well defined whenever $l < j-1$; otherwise, each of the above mentioned trees consists of its root only.

Then the number of elements of $\mathcal{E}^{(n)}_j \cap [0, \Theta^{(n)}_{j-1}(r)]$ on the leaves of $\mathfrak{T}_j^{(x_1,i|_l)}$, $l = 1, \ldots, j-2$, $j \ge 3$, is given by

$$\sum_{i_{l+2}=1}^{\tilde{G}^{(x_1,i|_{l+1})}} \cdots \sum_{i_j=1}^{\tilde{G}^{(x_1,i|_{j-1})}} 1 \tag{5.15}$$



Figure 4: Schematic representation of portions of $\cup_{i=1}^4 \mathcal{H}_i^{(n)}$ and the forest whose leaves are $\mathcal{E}_4^{(n)} \cap [0, \Theta_3^{(n)}(r)]$. Full points on horizontal dotted line represent $\mathcal{H}_1^{(n)} \cap [0, r]$. Successive generations of trees attached to each point of $\mathcal{H}_1^{(n)} \cap [0, r]$ represent $\mathcal{H}_2^{(n)} \cap [0, \Theta_1^{(n)}(r)]$, $\mathcal{H}_3^{(n)} \cap [0, \Theta_2^{(n)}(r)]$ and $\mathcal{H}_4^{(n)} \cap [0, \Theta_3^{(n)}(r)]$, respectively. Forest of extra marks and their descendants are shown in full lines and crosses. (Actual picture should look less regular, since the degrees of the vertices of the trees are independent random variables, which should be moreover large for large n.)

and their contribution to $\mu_j^{(n)}([0, \Theta_{j-1}^{(n)}(r)])$ amounts to

$$\sum_{i_{l+2}=1}^{\tilde{G}^{(x_1,i|_{l+1})}} \cdots \sum_{i_j=1}^{\tilde{G}^{(x_1,i|_{j-1})}} \gamma_j^{(n)}(x_1, U^{(x_1,i|_2)}, \ldots, U^{(x_1,i|_j)})\, T^{(x_1,i|_j)}, \tag{5.16}$$

where $\{T^{(x_1,i|_j)}\}$ are i.i.d. mean 1 exponential random variables, independent of all other random variables. So the size of $\mathcal{E}_j^{(n)} \cap [0, \Theta_{j-1}^{(n)}(r)]$ is given by

$$K_j^{(n)}(r) = \sum_{l=1}^{j-1} \sum_{x_1=1}^{M_1} \sum_{i_1=1}^{N_r^{(x_1,1)}} \sum_{i_2=1}^{G^{(x_1,i_1)}} \cdots \sum_{i_l=1}^{G^{(x_1,i|_{l-1})}} \sum_{i_{l+2}=1}^{\tilde{G}^{(x_1,i|_{l+1})}} \cdots \sum_{i_j=1}^{\tilde{G}^{(x_1,i|_{j-1})}} 1 \tag{5.17}$$

and its contribution to $\mu_j^{(n)}([0, \Theta_{j-1}^{(n)}(r)])$ amounts to

$$\mu_j^{(n)}(\mathcal{E}_j^{(n)} \cap [0, \Theta_{j-1}^{(n)}(r)]) =$$

$$\sum_{l=1}^{j-1} \sum_{x_1=1}^{M_1} \sum_{i_1=1}^{N_r^{(x_1,1)}} \sum_{i_2=1}^{G^{(x_1,i_1)}} \cdots \sum_{i_j=1}^{G^{(x_1,i|_{j-1})}} \sum_{i_{l+2}=1}^{\tilde{G}^{(x_1,i|_{l+1})}} \cdots \sum_{i_j=1}^{\tilde{G}^{(x_1,i|_{j-1})}} \gamma_j^{(n)}(x_1, U^{(x_1,i|_2)}, \ldots, U^{(x_1,i|_j)})\, T^{(x_1,i|_j)}, \tag{5.18}$$

where for $l = j - 1$, the expressions in (5.15-5.16) should be interpreted as 1 and

$$\gamma_j^{(n)}(x_1, U^{(x_1,i|_2)}, \ldots, U^{(x_1,i|_{j-1},1)})\, T^{(x_1,i|_{j-1},1)},$$

respectively, and for $j = 2$ (5.17-5.18) should be respectively interpreted as

$$K_2^{(n)}(r) = \sum_{x_1=1}^{M_1} \sum_{i_1=1}^{N_r^{(x_1,1)}} 1, \quad \mu_2^{(n)}(\mathcal{E}_2^{(n)} \cap [0, \Theta_1^{(n)}(r)]) = \sum_{x_1=1}^{M_1} \sum_{i_1=1}^{N_r^{(x_1,1)}} \gamma_2^{(n)}(x_1, U^{(x_1,i_1,1)})\, T^{(x_1,i_1,1)}. \tag{5.19}$$

We readily get from (5.19)

$$E(K_2^{(n)}(r)) = rM_1, \quad E(\mu_2^{(n)}(\mathcal{E}_2^{(n)} \cap [0, \Theta_1^{(n)}(r)])) = \frac{r}{M_2} \sum_{x|_2 \in \mathcal{M}|_2} \gamma_2^{(n)}(x|_2). \tag{5.20}$$

For $j \geq 3$, by conditioning on $N_r^{(x_1,1)}, G^{(x_1,i_1)}, \dots, G^{(x_1,i|_{j-2})}$, $U^{(x_1,i|_2)}, \dots, U^{(x_1,i|_{j-2})}$ (in the case of $j = 3$, $N_r^{(x_1,1)}, G^{(x_1,i_1)}$), and integrating on the remaining random variables, we get from (5.17)

$$\sum_{l=1}^{j-1} \sum_{x_1=1}^{M_1} \cdots \sum_{i_l=1}^{G^{(x_1,i|_{l-1})}} \sum_{i_{l+2}=1}^{\tilde{G}^{(x_1,i|_{l+1})}} \cdots \sum_{i_{j-1}=1}^{\tilde{G}^{(x_1,i|_{j-2})}} \frac{1}{M_{j-1}} \sum_{x_{j-1}=1}^{M_{j-1}} (1 + M_j\, \gamma_{j-1}^{(n)}(x_1, U^{(x_1,i_1)}, \dots, U^{(x_1,i|_{j-2})}, x_{j-1})).$$

Proceeding inductively, we find

$$E(K_j^{(n)}(r)) = \frac{r}{\prod_{p=2}^{j-1} M_p} \sum_{l=1}^{j-1} \sum_{x|_{j-1} \in \mathcal{M}|_{j-1}} \prod_{p=1}^{l-1} M_{p+1} \gamma_p^{(n)}(x|_p) \prod_{p=l+1}^{j-1} (1 + M_{p+1} \gamma_p^{(n)}(x|_p)). \tag{5.21}$$

Similarly,

$$E(\mu_j^{(n)}(\mathcal{E}_j^{(n)} \cap [0, \Theta_{j-1}^{(n)}(r)])) =$$
$$\frac{r}{\prod_{p=2}^{j} M_p} \sum_{l=1}^{j-1} \sum_{x|_j \in \mathcal{M}|_j} \gamma_j^{(n)}(x|_j) \prod_{p=1}^{l-1} M_{p+1} \gamma_p^{(n)}(x|_p) \prod_{p=l+1}^{j-1} (1 + M_{p+1} \gamma_p^{(n)}(x|_p)). \tag{5.22}$$

We are now ready to argue our claims.

a) Fix $j \in \{2, \dots, k\}, r > 0$ and $L > 0$. Then, using Jensen's inequality

$$\begin{aligned} P\left(\min\left\{\xi_j^{(n)}(s) : \, s \in \mathcal{E}_j^{(n)} \cap [0, \Theta_{j-1}^{(n)}(r)]\right\} > L\right) &= \textstyle\sum_{l=0}^{\infty} \left(1 - \frac{L}{M_j}\right)^l P\left(K_j^{(n)}(r) = l\right) \\ = E\left[\left(1 - \tfrac{L}{M_j}\right)^{K_j^{(n)}(r)}\right] \geq \left(1 - \tfrac{L}{M_j}\right)^{E(K_j^{(n)}(r))} &= \left\{\left(1 - \tfrac{L}{M_j}\right)^{M_j/L}\right\}^{L\left[E(K_j^{(n)}(r))/M_j\right]}. \end{aligned} \tag{5.23}$$

Using (5.21), we find that the expression within square brackets on the right hand side of (5.23) is the expression in (5.7), which goes to 0 as $n \to \infty$ by hypothesis. From (5.5), we have that the expression within curly brackets on the right hand side of (5.23) is bounded away from zero as $n \to \infty$. It immediately follows that the probability on the left hand side of (5.23) tends to 1 as $n \to \infty$, and part *a*) of Lemma 5.2 is established.

b) Given $\varepsilon > 0$, by Markov's inequality

$$P\left(\mu_j^{(n)}\left(\mathcal{E}_j^{(n)} \cap [0, \Theta_{j-1}^{(n)}(r)]\right) > \varepsilon\right) \leq \varepsilon^{-1} E\left(\mu_j^{(n)}\left(\mathcal{E}_j^{(n)} \cap [0, \Theta_{j-1}^{(n)}(r)]\right)\right) \tag{5.24}$$

and the result follows from (5.22) and (5.8).

□

Remark 5.3. *If $X_{k-1}^{(n)} \to X_{k-1}$ a.s. as $n \to \infty$ in the Skorohod space, then, by Proposition 5.2 in Chapter 3 of [24] (page 118), we have that*

$$\lim_{n\to\infty} X_{k-1}^{(n)}(s) = \lim_{n\to\infty} X_{k-1}^{(n)}(s-) = X_{k-1}(s) \tag{5.25}$$

for all $s \geq 0$ which is a continuity point of X_{k-1}.

Lemma 5.4. *Assume that the induction hypotheses (5.10-5.11) hold and let $r \in [0,\infty)$ be fixed. Then $\Gamma_k^{(n)}(r) \to \Gamma_k(r)$ in probability as $n \to \infty$.*

Proof. The strategy is to separate the contribution of the extra marks and Poissonian marks with large labels from the remaining contributions. The convergence of the remainder, main (as it turns out) contributions to the corresponding infinite volume ones follows readily from the first part of (5.6), since there is only a fixed finite number of contributions involved. The negligibility of the total contribution of extra marks was established in Lemma 5.2 above, so we are left with establishing that of the total contribution of high label marks. Details follow.

Let $\Psi_k^{(n)}(r) = \mu_k^{(n)}\Big((\mathcal{S}_k^{(n)} \setminus \mathcal{E}_k^{(n)}) \cap [0,r]\Big)$. Then

$$|\Gamma_k(r) - \Gamma_k^{(n)}(r)| \leq |\Gamma_k(r) - \Psi_k^{(n)}(r)| + \mu_k^{(n)}(\mathcal{E}_k^{(n)} \cap [0,r]). \tag{5.26}$$

By Lemma 5.2.b, the second term goes to 0 in probability as $n \to \infty$. We will argue that so does the first one. In preparation for that, let us take, for given $\varepsilon > 0$, $m_1 \in \mathbb{N}_*$ such that

$$\sum_{x_1 > m_1} \sum_{x|^2 \in \mathbb{N}_*^{k-1}} \bar{\gamma}_k(x|_k) \leq \varepsilon/k, \tag{5.27}$$

(recall notation introduced around (2.2) above). Proceeding inductively, with $m_1, \ldots, m_{j-1}$, $2 \leq j \leq k-1$, fixed, choose m_j such that

$$\sum_{x_1=1}^{m_1} \cdots \sum_{x_{j-1}=1}^{m_{j-1}} \sum_{x_j > m_j} \sum_{x|^{j+1} \in \mathbb{N}_*^{j+1}}^{\infty} \bar{\gamma}_k(x|_k) \leq \varepsilon/k, \tag{5.28}$$

and with $m_1, \ldots, m_{k-1}$ fixed, choose m_k such that

$$\sum_{x_1=1}^{m_1} \cdots \sum_{x_{k-1}=1}^{m_{k-1}} \sum_{x_k > m_k}^{\infty} \bar{\gamma}_k(x|_k) \leq \varepsilon/k. \tag{5.29}$$

This procedure is well defined by (4.2).

Going back to the first term on the right of (5.26), we have that

$$|\Gamma_k(r) - \Psi_k^{(n)}(r)| \leq \left| \sum_{s \in \tilde{\mathcal{S}}_k^{(m_k)} \cap [0,r]} \left\{ \gamma_k(X_{k-1}(s), \xi_k(s)) - \gamma_k^{(n)}(X_{k-1}^{(n)}(s), \xi_k(s)) \right\} T_s^{(k)} \right| \tag{5.30}$$

$$+ \sum_{s \in (\mathcal{S}_k \setminus \tilde{\mathcal{S}}_k^{(m_k)}) \cap [0,r]} \gamma_k(X_{k-1}(s), \xi_k(s))\, T_s^{(k)} \tag{5.31}$$

$$+\sum_{s\in(\mathcal{S}_k^{(n)}\setminus\tilde{\mathcal{S}}_k^{(m_k)})\cap[0,r]}\gamma_k^{(n)}(X_{k-1}^{(n)}(s),\xi_k(s))\,T_s^{(k)}. \tag{5.32}$$

The expression on the right hand side of (5.30) converges to 0 in probability as n increases because it is a finite sum, and from the first part of (5.6), and since $X_{k-1}^{(n)}(s)=X_{k-1}(s)$ for all $s\in\tilde{\mathcal{S}}_k^{(m_k)}\cap[0,r]$ for all large enough n almost surely, as follows from Remark 5.3 above.

Let B and C denote the expressions in (5.31) and (5.32), respectively.

To analyse B, we start by taking, for given $\eta>0$, r_0' such that

$$P(\Theta_{k-1}(r_0')>r)\geq 1-\eta. \tag{5.33}$$

That this is allowed follows from Remark 4.6. Now letting

$$\mathcal{A}_0=(\mathcal{S}_k\setminus\tilde{\mathcal{S}}_k^{(m_k)})\cap[0,\Theta_{k-1}(r_0')],$$

we define

$$\begin{aligned}\mathcal{A}_1 &= \{s\in\mathcal{A}_0:\,X_{k-1,1}(s)>m_1\},\\ \mathcal{A}_2 &= \{s\in\mathcal{A}_0:\,X_{k-1,1}(s)\leq m_1 \text{ and } X_{k-1,2}>m_2\}\\ &\vdots\\ \mathcal{A}_{k-1} &= \{s\in\mathcal{A}_0:\,X_{k-1,1}(s)\leq m_1,\ldots,X_{k-1,k-1}\leq m_{k-1}\}.\end{aligned} \tag{5.34}$$

Then, outside an event of probability at most η, we have that

$$B\leq\sum_{i=1}^{k-1}\sum_{s\in\mathcal{A}_i}\gamma_k(X_{k-1}(s),\xi_k(s))\,T_s^{(k)}, \tag{5.35}$$

and following the same arguments used to establish (4.11), and using (5.27-5.29), we conclude that

$$E\left(\sum_{i=1}^{k-1}\sum_{s\in\mathcal{A}_i}\gamma_k(X_{k-1}(s),\xi_k(s))\,T_s^{(k)}\right)=$$
$$r_0'\left[\sum_{x_1>m_1}\sum_{x|^2\in\mathbb{N}_*^{k-1}}\bar{\gamma}_k(x|_k)+\sum_{x_1=1}^{m_1}\sum_{x_2>m_2}\sum_{x|^3\in\mathbb{N}_*^{k-2}}\bar{\gamma}_k(x|_k)+\ldots+\sum_{x_1=1}^{m_1}\cdots\sum_{x_{k-1}=1}^{m_{k-1}}\sum_{x_k>m_k}^{\infty}\bar{\gamma}_k(x|_k)\right]$$
$$\leq r_0'\,\varepsilon. \tag{5.36}$$

This shows that $B\to 0$ in probability as $\epsilon\to 0$, since η is arbitrary.

The analysis of C is similar, with the dependence on n as a distinctive aspect. From induction hypothesis (5.11) and Remark 4.6, given $\eta>0$, there exists r_0 such that for all n suficiently large

$$P(\Theta_{k-1}^{(n)}(r_0)>r)\geq 1-\eta \tag{5.37}$$

(recall (5.13)). With such r_0 and the above choice of $m_1,\ldots,m_k$, define

$$\mathcal{A}_0^{(n)}=\left(\mathcal{S}_k^{(n)}\setminus\mathcal{S}_k^{(m_k)}\right)\cap[0,\Theta_{k-1}^{(n)}(r_0)],$$

$$\begin{aligned}
\mathcal{A}_1^{(n)} &= \{s \in \mathcal{A}_0^{(n)} : X_{k-1,1}^{(n)}(s) > m_1\}, \\
\mathcal{A}_2^{(n)} &= \{s \in \mathcal{A}_0^{(n)} : X_{k-1,1}^{(n)}(s) \le m_1 \text{ and } X_{k-1,2}^{(n)} > m_2\} \\
&\vdots \\
\mathcal{A}_k^{(n)} &= \{s \in \mathcal{A}_0^{(n)} : X_{k-1,1}^{(n)}(s) \le m_1, \dots, X_{k-1,k-1}^{(n)} \le m_{k-1}\}.
\end{aligned} \tag{5.38}$$

Then

$$P\left(C \le \sum_{i=1}^{k-1} \sum_{s \in \mathcal{A}_i^{(n)}} \gamma_k^{(n)}(X_{k-1}^{(n)}(s), \xi_k(s)) T_s^{(k)} \right) \ge 1 - \eta \tag{5.39}$$

for all n large enough. By (5.6), we may take n suficiently large such that

$$\begin{gathered}
\left| \sum_{x_1 > m_1}^{M_1} \sum_{x|^2 \in \mathcal{M}|^2} \bar{\gamma}_k^{(n)}(x|_k) - \sum_{x_1 > m_1} \sum_{x|^2 \in \mathbb{N}_*^{k-1}} \bar{\gamma}_k(x|_k) \right| \le \varepsilon/k, \\
\left| \sum_{x_1=1}^{m_1} \sum_{x_2=m_2+1}^{M_2} \sum_{x|^3 \in \mathcal{M}|^3} \bar{\gamma}_k^{(n)}(x|_k) - \sum_{x_1=1}^{m_1} \sum_{x_2 > m_2} \sum_{x|^3 \in \mathbb{N}_*^{k-2}} \bar{\gamma}_k(x|_k) \right| \le \varepsilon/k, \\
\vdots \\
\left| \sum_{x_1=1}^{m_1} \cdots \sum_{x_{k-1}=1}^{m_{k-1}} \sum_{x_k=m_k+1}^{M_k} \bar{\gamma}_k^{(n)}(x|_k) - \sum_{x_1=1}^{m_1} \cdots \sum_{x_{k-1}=1}^{m_{k-1}} \sum_{x_k > m_k} \bar{\gamma}_k(x|_k) \right| \le \varepsilon/k.
\end{gathered} \tag{5.40}$$

Following the same arguments used to establish (4.11), and using (5.27-5.29) and (5.40), we get

$$\begin{gathered}
E\left(\sum_{i=1}^{k-1} \sum_{s \in \mathcal{A}_i^{(n)}} \gamma_k^{(n)}(X_{k-1}^{(n)}(s), \xi_k(s))\, T_s^{(k)} \right) = \\
r_0' \left[\sum_{x_1=m_1+1}^{M_1} \sum_{x|^2 \in \mathcal{M}|^2} \bar{\gamma}_k^{(n)}(x|_k) + \sum_{x_1=1}^{m_1} \sum_{x_2=m_2+1}^{M_2} \sum_{x|^3 \in \mathcal{M}|^3} \bar{\gamma}_k^{(n)}(x|_k) + \dots + \sum_{x_1=1}^{m_1} \cdots \sum_{x_{k-1}=1}^{m_{k-1}} \sum_{x_k=m_k+1}^{M_k} \bar{\gamma}_k^{(n)}(x|_k) \right] \\
\le (r_0 + r_0')\, \varepsilon.
\end{gathered}$$

This shows that $C \to 0$ in probability as we first take $n \to \infty$ and then $\epsilon \to 0$, since η is arbitrary, thus completing the proof.

□

Corollary 5.5. *The result of Lemma 5.4 still holds if we replace r on the left hand side by r_n with $r_n \to r$ as $n \to \infty$, with (r_n) a deterministic sequence.*

Proof. Let us write

$$|\Gamma_k^{(n)}(r_n) - \Gamma_k(r)| \le |\Gamma_k^{(n)}(r_n) - \Gamma_k^{(n)}(r)| + |\Gamma_k^{(n)}(r) - \Gamma_k(r)|. \tag{5.41}$$

Using the hypothsis and the monotonicity of $\Gamma_k^{(n)}$, given $\delta > 0$, we have that the first term on the right hand side is bounded above by $\Gamma_k^{(n)}(r_\delta) - \Gamma_k^{(n)}(r-\delta)$ for all n large enough, which is in turn bounded above by

$$\Gamma_k(r+\delta) - \Gamma_k(r-\delta) + |\Gamma_k^{(n)}(r+\delta) - \Gamma_k(r+\delta)| + |\Gamma_k^{(n)}(r-\delta) - \Gamma_k(r-\delta)| \tag{5.42}$$

Let $\eta > 0$ now be given. By Lemma 5.4, and using (5.41-5.42), we find that

$$\limsup_{n\to\infty} P(|\Gamma_k^{(n)}(r_n) - \Gamma_k(r)| > \eta) \le P(\Gamma_k(r+\delta) - \Gamma_k(r-\delta) > \eta/4), \tag{5.43}$$

and the result follows from r being almost surely a continuity point of Γ_k (see Remarks 4.1 and 4.3), since δ is arbitrary.

□

Remark 5.6. *The same argument of course works in the case* (r_n, r) *are random and independent of* $(\Gamma_k^{(n)}, \Gamma_k)$ *and* $r_n \to r$ *almost surely as* $n \to \infty$. *This can be applied to establish that under the assumption of Lemma 5.4, we have that*

$$\Theta_k^{(n)}(T) \to \Theta_k(T) \text{ as } n \to \infty, \tag{5.44}$$

in probability for every $T \ge 0$.

The next result is a finite volume version of Lemma 4.8 in the above section.

Lemma 5.7. *Given* $k \ge 2$, $n \ge 1$, *let* $X_k^{(n)}$ *be a trap model on* $\mathbb{T}_k^{(n)}$. *Suppose that the assumptions of Theorem 5.1 and the induction hypotheses (5.10-5.11) all hold. Then given* $m \ge 1$ *and* $\epsilon > 0$, *there exists* $\tilde m = \tilde m^{(k)} \ge 1$ *such that the event*

$$\begin{aligned} A_m^{(n)} \; = \; \{ &\text{if } X_{k,i}^{(n)}(t) > \tilde m \text{ for some } i = 1,\dots,k-1 \text{ and } t \in [0,T], \\ &\text{then } X_{k,j}^{(n)}(t) > m \text{ for } j = i+1,\dots,k\} \end{aligned} \tag{5.45}$$

has probability bounded below by $1-\epsilon$ *for all* n *large enough.*

Proof. We argue similarly as in the proof of Lemma 4.8, except that statements hold *with high probability*, instead of almost surely.

By Remark 5.6 and the a.s. finiteness of $\Theta_j(r)$ for every $1 \le j \le k$ and $r \ge 0$, we may choose $T' > 0$ be such that $\Theta_k^{(n)}(T') > T$ with probability at least $1-\epsilon/4$ uniformly in n. Now for $m \in \mathbb{N}_*$ and $j = 1,\dots,k$, let $\tilde{\mathcal{S}}_j^{(m)}$ be as in (4.14) above. For fixed $\ell \in \mathbb{N}_*$, let $\mathcal{T}_{kj}^{(n)}(\ell)$ denote the set of times up to $\Theta_{k-1}^{(n)}(T')$ spent by $X_{k-1,j}^{(n)}$ above ℓ, $j = 1,\dots,k-1$. Analogously as for the infinite volume case (see (4.15)), we may check that the expected length of $\mathcal{T}_{kj}^{(n)}(\ell)$ equals

$$T' \sum_{x_1=1}^{M_1} \cdots \sum_{x_j=\ell+1}^{M_j} \cdots \sum_{x_{k-1}=1}^{M_{k-1}} \gamma_1^{(n)}(x|_1)\dots\gamma_{k-1}^{(n)}(x|_{k-1}) \tag{5.46}$$

plus the contribution of the extra marks and their descendants. It follows from (5.6) that the $\limsup_{n\to\infty}$ of the expression in (5.46) vanishes as $\ell \to \infty$. By Lemma 5.2.b, the contribution of the extra marks and their descendants vanishes in probability as $n \to \infty$. It then follows from elementary properties of Poisson processes, that

$$\{\cup_{j=1}^{k-1}\mathcal{T}_{kj}^{(n)}(\ell)\} \cap \tilde{\mathcal{S}}_k^{(m)} = \emptyset \tag{5.47}$$

outside an event whose probability is bounded above by $\epsilon/4$ for all ℓ, n large enough. This statement is about Poissonian marks; but it also holds for extra marks by Lemma 5.2.*a*.

So, given $m \in \mathbb{N}_*$ and $\epsilon > 0$, we find $\hat{m}^{(k)}$ such that outside an event of probability smaller than $\epsilon/2$ for all n large enough, if $X_{k,i}^{(n)}(t) > \hat{m}^{(k)}$ for any $t \leq T$, then $X_{k,k}^{(n)}(t) > m$. This in particular establishes the claim for $k = 2$ by the choice $\tilde{m}^{(2)} = \hat{m}^{(2)}$. Let us assume that the claim is established for $k-1$. Then substituting $\epsilon/4$ as ϵ, T'' such that $P(\Theta_{k-1}^{(n)}(T') \leq T'') > 1 - \epsilon/4$ for all large enough n as T, and $\hat{m}^{(k)} \vee m$ as m in that claim, and taking the choice $\tilde{m}^{(k)} = \tilde{m}^{(k-1)} \vee \hat{m}^{(k)}$, $i = 1, \ldots, k-1$, we find that it satisfies the claim.

$\square$

5.3 End of proof of Theorem 5.1

For $j = 1, \ldots, k$, $n \geq 1$, let $X_j^{(n)} \sim TM(\mathbb{T}_k^{(n)}, \underline{\gamma_j^{(n)}})$ and $X_j \sim K_j(\mathbb{T}_k, \underline{\gamma_j})$, and, for $k \geq 2$ fixed, suppose that $X_{k-1}^{(n)} \to X_{k-1}$ in probability as $n \to \infty$. We will/may then inductively suppose that

$$X_{k-1}^{(n')} \to X_{k-1} \text{ a.s. as } n' \to \infty, \tag{5.48}$$

for a subsequence (n'). We will fix $\epsilon > 0$, $T > 0$ and $m \geq 1$ and choose T' and $\tilde{m}$ such that outside an event $\mathcal{E} = \mathcal{E}_{n'}$ of probability at most $\epsilon/2$ for all n' large enough we have that the conclusions of Lemma 4.8 and 5.7 hold, and also that $\Theta_{k-1}(T') \wedge \Theta_{k-1}^{(n')}(T') > T$. We will also assume that $\tilde{m} \geq m$, and that the claims of Lemma 5.2 hold almost surely over (n').

On the way to showing the validity of (5.48) with k replacing $k-1$ (in probability), we now proceed to define appropriate time distortions $\lambda^{(n')}$ (see discussion on the Skorohod metric at the beginning of Subsection 5.1). Let us start by considering the constancy intervals of $X_{k-1,1}$ in $[0, \Theta_{k-1}(T'))$ with $X_{k-1,1} \leq \tilde{m}$. These are defined to be the *rank-$\tilde{m}$ constancy intervals of the level* 1 *for* X_{k-1}. Proceeding inductively, given $2 \leq \ell \leq k-1$, for each rank-$\tilde{m}$ constancy interval I of level $\ell - 1$, we consider the constancy intervals of $X_{k-1,\ell}$ inside I such that $X_{k-1,\ell} \leq \tilde{m}$. The collection of all such intervals obtained from all the rank-$\tilde{m}$ constancy intervals of level $\ell - 1$ for X_{k-1} form the set of rank-$\tilde{m}$ constancy intervals of level ℓ for X_{k-1}.

Let $a_1, \ldots, a_{2L}$ denote the collection of all endpoints of all the rank-$\tilde{m}$ constancy intervals of level i for X_{k-1}, $i = 1, \ldots, k-1$, in increasing order, and let $b_1, \ldots, b_{2J}$ denote the collection of all endpoints of all the rank-$\tilde{m}$ constancy intervals of level $k-1$ for X_{k-1} in increasing order. See Figure 5.

Let us also consider rank-$\tilde{m}$ constancy intervals of level i for $X_{k-1}^{(n')}$, with the paralell definition to the one above. By the assumption that Lemma 5.2 holds almost surely over (n'), and for n' large enough, there is one-to-one correspondence of the $a_1^{(n')}, \ldots, a_{2L^{(n')}}^{(n')}$ and $a_1, \ldots, a_{2L}$, with $L^{(n')} = L$ for all large n' and $a_i^{(n')}$ corresponding to a_i, and from (5.48),

$$a_i^{(n')} \to a_i \tag{5.49}$$



Figure 5: Depiction of objects appearing in the argument for the proof of Theorem 5.1 with $k = 3$. Rank-$\tilde{m}$ constancy intervals of the first level ($[a_1, a_6)$ and $[a_7, a_{14})$) and rank-$\tilde{m}$ constancy intervals of the second level ($[a_2, a_3)$, $[a_4, a_5)$, $[a_8, a_9)$, $[a_{10}, a_{11})$ and $[a_{12}, a_{13})$) for X_2 appear on the x-axis. Correspondingly on the y-axis, we have rank-$\tilde{m}$ constancy intervals of the first level ($[A_1, A_6)$ and another one whose endpoints are not named in the picture), rank-$\tilde{m}$ constancy intervals of the second level ($[A_2, A_3)$, $[A_4, A_5)$, and other ones whose endpoints are not named in the picture), and rank-$\tilde{m}$ constancy intervals of the third level ($[A_1^-, A_1^+)$, $[A_2^-, A_2^+)$, $[A_3^-, A_3^+)$, and other ones whose endpoints are not named in the picture) for X_3. Some of the correspondences between the axes are indicated by dotted lines. We have also $b_1 = a_2$, $b_2 = a_3$, $b_3 = a_4$, $b_4 = a_5$, $b_5 = a_8$, $b_6 = a_9$, $b_7 = a_{10}$, $b_8 = a_{11}$, $b_9 = a_{12}$ and $b_{10} = a_{13}$. This picture is also good for $X_3^{(n)}$, with n large, and with all endpoint labels having superscripts "(n)".

almost surely as $n' \to \infty$ for every $i = 1, \dots, 2L$.

Let now $A_i = \Gamma_k(a_i)$ and $A_i^{(n')} = \Gamma_k^{(n')}(a_i^{(n')})$, $i = 1, \dots, 2L$. See Figure 5. It follows from Lemma 5.4 that

$$A_i^{(n')} \to A_i \tag{5.50}$$

in probability as $n' \to \infty$, and we may assume a.s. convergence by taking a subsequence.

Now let $\{s'_1, \dots, s'_Q\}$ be the enumeration in increasing order of $\cup_{i=1}^J \{\tilde{\mathcal{S}}_k^{(m)} \cap [b_{2i-1}, b_{2i})\}$, and let $A_i^- = \Gamma_k(s'_i-)$ and $A_i^+ = \Gamma_k(s'_i)$. See Figure 5. We note that the intervals $[A_i^-, A_i^+)$, $i = 1, \dots, Q$ are the rank-$\tilde{m}$ constancy intervals of level k for X_k, with $m_k^{(k)} = m$, whereas A_i, $i = 1, \dots, 2L$, are the endpoints of all the rank-$\tilde{m}$ constancy intervals of level i for X_k, $i = 1, \dots, k-1$.

We remark at this point that under our assumptions so far, we have that for all $i = 1, \dots, J$

$$\tilde{\mathcal{S}}_k^{(m)} \cap [b_{2i-1}, b_{2i}) = \tilde{\mathcal{S}}_k^{(m)} \cap [b_{2i-1}^{(n')}, b_{2i}^{(n')}) \tag{5.51}$$

almost surely for all n' large enough, where $b_i^{(n')} = a_j^{(n')}$ such that $a_j = b_i$.

Let $A_i^-(n') = \Gamma_k^{(n')}(s'_i-)$, $A_i^+(n') = \Gamma_k^{(n')}(s'_i)$. Then for all large enough n', $[A_i^-(n'), A_i^+(n'))$, $i = 1, \dots, Q$ are the rank-$\tilde{m}$ constancy intervals of level k for $X_k^{(n')}$, with $m_k^{(k)} = m$, whereas $A_i^{(n')}$, $i = 1, \dots, 2L$, are the endpoints of all the rank-$\tilde{m}$ constancy intervals of level i for $X_k^{(n')}$, $i = 1, \dots, k-1$.

Let us now argue that

$$A_i^{\pm}(n') \to A_i^{\pm} \tag{5.52}$$

in probability as $n' \to \infty$ (and again we may assume a.s. convergence by taking a subsequence). Indeed it is enough to first note that almost surely $A_i^-(n') = \tilde{\Gamma}_k^{(n')}(s'_i)$, $A_i^- = \tilde{\Gamma}_k(s'_i)$, $A_i^+(n') = \tilde{\Gamma}_k^{(n')}(s'_i) + \gamma_k^{(n)}(X_{k-1}^{(n')}(s'_i), \xi_k(s'_i))$ and $A_i^+ = \tilde{\Gamma}_k(s'_i) + \gamma_k(X_{k-1}(s'_i), \xi_k(s'_i))$, where $\tilde{\Gamma}_k^{(n')}$ and $\tilde{\Gamma}_k$ are obtained from $\tilde{\mu}_k^{(n')}$ and $\tilde{\mu}_k$ as $\Gamma_k^{(n')}$ and $\Gamma_k^{(n')}$ are obtained from $\mu_k^{(n')}$ and μ_k, respectively, where $\tilde{\mu}_k^{(n')} = \mu_k^{(n')}$ and $\tilde{\mu}_k = \mu_k$ everywhere except at $\{s'_i\}$, where $\tilde{\mu}_k^{(n')}$ and $\tilde{\mu}_k$ both vanish. By the same arguments above we get $\tilde{\Gamma}_k^{(n')}(s'_i) \to \tilde{\Gamma}_k(s'_i)$ in probability, and the result follows upon noticing that $X_{k-1}^{(n')}(s'_i) = X_{k-1}(s'_i)$ for all large enough n' and using (5.6).

We are now ready to define our time distortion. Let $\lambda^{(n')} : [0, \infty) \to [0, \infty)$ be such that

$$\lambda^{(n')}(A_i) = A_i^{(n')}, \quad \lambda^{(n')}(A_i^-) = A_i^-(n'), \quad \lambda^{(n')}(A_i^+) = A_i^+(n'), \tag{5.53}$$

and make it linear between successive points of $\mathcal{A} := \{A_i,\, i = 1, \dots, 2L;\, A_j^-, A_j^+,\, j = 1, \dots, Q\}$, and linear with inclination 1 from $\max \mathcal{A}$ on. Then $\lambda^{(n')}$ is almost surely well defined for all large enough n', and one readily checks that condition (5.53) implies that $\lambda^{(n')}$ maps rank-$\tilde{m}$ constancy intervals of level i for X_k to the corresponding rank-$\tilde{m}$ constancy intervals of level i for $X_k^{(n')}$, $i = 1, \dots, k$, given by the coupling. In particular, $X_{k,j}(\lambda^{(n')}(\cdot)) = X_{k,j}^{(n')}(\cdot)$, $j = 1, \dots i$, on those respective intervals. From the assumptions of the paragraph of (5.48), we then have that outside $\mathcal{E}$

$$\sup_{0 \le u \le T} \rho(X_k^{(n')}, X_k, \lambda^{(n')}, u) \le 1/m \tag{5.54}$$

and by (5.50) and (5.52) and our construction and assumptions it follows that

$$\phi(\lambda^{(n')}) \to 0 \tag{5.55}$$

as $n' \to \infty$ almost surely.

It follows from all of the above that for every fixed $\epsilon, T > 0$ and $m \in \mathbb{N}_*$ we may find a subsequence (n') such that

$$P\left(\rho(X_k^{(n')}, X_k) > \frac{1}{m} + e^{-T}\right) \le \epsilon, \tag{5.56}$$

for all n' large enough, so we have that $X_k^{(n')} \to X_k$ in probability, and this readily implies the claim of Theorem 5.1.

6 Scaling limit of the GREM-like trap model

In this section we apply Theorem 5.1 to obtain a scaling limit of the GREM-like trap model in the fine tuning regime and extreme time scale. We first define these terms.

For $j = 1, \dots, k$, let $\tau_j := \{\tau_j(x|_j);\ x|_j \in \mathcal{M}|_j\}$ be an i.i.d. family of random variables in the domain of attraction of an α_j-stable law. We suppose

$$0 < \alpha_1 < \cdots < \alpha_k < 1. \tag{6.1}$$

For $j = 1, \dots, k$, we will relabel τ_j obtaining $\tau_j^{(n)} = \{\tau_j^{(n)}(x|_j);\ x|_j \in \mathcal{M}|_j\}$, so that, for every $(x|_{j-1}) \in \mathcal{M}|_{j-1}$, $\{\tau_j^{(n)}(x|_j);\ x_j \in \mathcal{M}_j\}$ are the decreasing order statistics of $\{\tau_j(x|_j);\ x_j \in \mathcal{M}_j\}$.

For $j = 1, \dots, k$, $n \ge 1$, let

$$c_j^{(n)} = (G_j^{-1}(M_j^{-1}))^{-1}, \tag{6.2}$$

where G_j^{-1} is the (generalized) inverse of $G_j : [0, \infty) \to (0, 1]$ such that $G_j(t) = P(\tau_j(x|_j) > t)$.

Then, making

$$\gamma_j^{(n)}(x|_j) = c_j^{(n)} \tau_j^{(n)}(x|_j),\ x|_j \in \mathcal{M}_j, \tag{6.3}$$

we have that for all $j = 1, \dots, k$ and $x|_{j-1} \in \mathbb{N}_*^{j-1}$, the sequence of point processes

$$\sum_{x_j \in \mathcal{M}_j} \mathbf{1}_{\gamma_j^{(n)}(x|_j)} \tag{6.4}$$

converges in distribution as $n \to \infty$ to independent Poisson point processes

$$\sum_{x_j \in \mathbb{N}_*} \mathbf{1}_{\gamma_j(x|_j)} \tag{6.5}$$

with intensity measure given by the density $y^{-\alpha_j - 1}$, $y > 0$. We assume that for all $j = 1, \dots, k$ and $x|_{j-1} \in \mathbb{N}_*^{j-1}$ $\gamma_j(x|_j)$ is non increasing in x_j.

The *fine tuning* regime mentioned above and at the introduction corresponds to choosing $M_1 = n$ and

$$M_{j+1} = \left\lfloor 1/c_j^{(n)} \right\rfloor, \quad j = 1, \dots, k-1. \tag{6.6}$$

We will also assume this choice from now.

Let

$$\tilde{\gamma}_j^{(n)}(x|_j) = \begin{cases} \gamma_j^{(n)}(x|_j), & \text{if } j = 1, \dots, k-1, \\ \tau_k^{(n)}(x|_k), & \text{if } j = k. \end{cases} \tag{6.7}$$

and let $\tilde{X}_k^{(n)} \sim TM(\mathbb{T}_k^{(n)}; \underline{\tilde{\gamma}_k^{(n)}})$.

Remark 6.1. *One may readily check from Lemma 3.7 that, in terms of the coin tossing description,* $\tilde{X}_k^{(n)} \sim TM(\mathbb{T}_k^{(n)}; \tau_k^{(n)}, (p_j^{(n)})_{j=1}^{k-1})$*, where for* $j = 1, \dots, k-1$ *and* $x|_j \in \mathcal{M}|_j$

$$p_j^{(n)}(x|_j) = \frac{1}{1 + \tau_j^{(n)}(x|_j)}. \tag{6.8}$$

With this description, and general finite $M_1, \dots, M_k$ *(not necessarily satisfying (6.6)), we call* $\tilde{X}_k^{(n)}$ *the GREM-like trap model on* $\mathbb{T}_k^{(n)}$ *with parameters* $\tau_j^{(n)}(x|_j)$*,* $j = 1, \dots, k$*,* $x|_j \in \mathcal{M}|_j$*.*

Let us speed up $\tilde{X}_k^{(n)}$ by $c_k^{(n)}$, namely, let

$$X_k^{(n)} = \tilde{X}_k^{(n)}(t/c_k^{(n)}), \quad t \geq 0. \tag{6.9}$$

This corresponds to the extreme time scale mentioned above and at the introduction. One may readily check that $X_k^{(n)} \sim TM(\mathbb{T}_k^{(n)}; \underline{\gamma_k^{(n)}})$. Let $X_k \sim K(\mathbb{T}_k; \underline{\gamma_k})$.

Theorem 6.2. *Let* $X_k^{(n)}$ *and* X_k *be as above. Then*

$$(X_k^{(n)}, \underline{\gamma_k^{(n)}}) \Rightarrow (X_k, \underline{\gamma_k}), \tag{6.10}$$

where $\Rightarrow$ *means weak convergence in the product of Skorohod space with the space of finite measures in* $\mathbb{N}_*^k$ *equipped with the topology of weak convergence.*

Proof. The strategy will be to work with a coupled version of $(\underline{\gamma_k^{(n)}}, \underline{\gamma_k})$, which we will call $(\underline{\hat{\gamma}_k^{(n)}}, \underline{\hat{\gamma}_k})$, such that almost surely for every $j = 1, \dots, k$ and $x|_j \in \mathbb{N}_*^j$

$$\hat{\gamma}_j^{(n)}(x|_j) \to \hat{\gamma}_j(x|_j) \ \text{ as } \ n \to \infty, \tag{6.11}$$

and then verifying the remaining conditions of Theorem 5.1.

For the coupling, we use the construction of [10], Section 6, which we describe briefly, guiding the reader to that reference for more details.

Let $E_j(x|_j)$, $x|_j \in \mathbb{N}_*^j$, $j = 1, \dots, k$ be independent mean one exponential random variables, and, for $x|_{j-1} \in \mathbb{N}_*^{j-1}$ make

$$S_j(x|_j) = \sum_{i=1}^{x_j} E_j(x|_{j-1}, i), \tag{6.12}$$

where $x|_0$ is a void symbol. Let now

$$\hat{\gamma}_j^{(n)}(x|_j) = c_j^{(n)} G_j^{-1}\left(\frac{S_j(x|_j)}{S_j(x|_{j-1}, M_j + 1)} \right); \tag{6.13}$$

$$\hat{\gamma}_j(x|_j) = S_j(x|_j)^{-1/\alpha_j}. \tag{6.14}$$

From an elementary large deviation estimate, we may assume that

$$S_j(x|_{j-1}, M_j + 1) \leq 2M_j \tag{6.15}$$

for all $x|_{j-1} \in \mathcal{M}|_{j-1}$ and n sufficiently large almost surely.

Then $\hat\gamma_j^{(n)}(x|_{j-1})$ and $\hat\gamma_j(x|_{j-1})$ are versions of $\gamma_j^{(n)}(x|_{j-1})$ and $\gamma_j(x|_{j-1})$, respectively [25]. Proposition 6.3 and Lemma 6.4 of [10] immediately implies (6.11), and also that almost surely for every $j = 1, \dots, k$ and $x|_{j-1} \in \mathbb{N}_*^{j-1}$

$$\sum_{x_j \in \mathcal{M}_j} \hat\gamma_j^{(n)}(x|_j) \to \sum_{x_j \in \mathbb{N}_*} \hat\gamma_j(x|_j) \text{ as } n \to \infty. \tag{6.16}$$

The a.s. validity of the first part of (5.6) for all j, x, as well as that of the second part for $k = 1$, follows immediately.

In order to get the a.s. validity of the second part of (5.6) for general k, we argue as follows. We may suppose by induction that it holds for $k-1$. We first write the sum in the second part of (5.6) more explicitly as follows:

$$\sum_{x_1} \hat\gamma_1^{(n)}(x_1) \cdots \sum_{x_k} \hat\gamma_k^{(n)}(x|_k), \tag{6.17}$$

and break each of the k sums above (following the strategy of [10]; see proof of Proposition 6.3) in three parts, so that the j-th sum is written as

$$\sum_{x_j}^{(1)} + \sum_{x_j}^{(2)} + \sum_{x_j}^{(3)}, \tag{6.18}$$

where given $\delta_j \in (0,1)$, the first sum is over $x|_j$ such that $\hat\gamma_k(x|_j) > \delta_j$, the second sum is over $x|_j$ such that $M_j^{-1/\alpha_j} < \hat\gamma_k(x|_j) \leq \delta_j$, and the third sum is over $x|_j$ such that $\hat\gamma_k(x|_j) \leq M_j^{-1/\alpha_j}$.

It follows from Lemma 6.4 of [10] that

$$\sum_{x_1}^{(1)} \hat\gamma_1^{(n)}(x_1) \cdots \sum_{x_k}^{(1)} \hat\gamma_k^{(n)}(x|_k) \to \sum_{x_1}^{(1)} \hat\gamma_1(x_1) \cdots \sum_{x_k}^{(1)} \hat\gamma_k(x|_k) \tag{6.19}$$

as $n \to \infty$ almost surely, since these are sums over a fixed bounded set of terms. We will show that

$$\limsup_{\delta_1,\dots,\delta_k \to 0} \limsup_{n \to \infty} \sum_{x_1}^{(i_1)} \hat\gamma_1^{(n)}(x_1) \cdots \sum_{x_k}^{(i_k)} \hat\gamma_{i_k}^{(n)}(x|_k) = 0 \tag{6.20}$$

almost surely, for all $(i_1, \dots, i_k) \in \{1,2,3\}^k \setminus \{(1,\dots,1)\}$. Since, again, $\sum_{x_j}^{(1)}$ are sums over a fixed bounded set of terms, and using the induction hypothesis, it is enough to consider sums

$$\sum_{x_1}^{(i_1)} \hat\gamma_1^{(n)}(x_1) \cdots \sum_{x_k}^{(i_k)} \hat\gamma_k^{(n)}(x|_k) \tag{6.21}$$

with $i_1 \in \{2,3\}$.

Let us first consider the case where $i_1 = 2$ and $i_j \in \{1,2\}$ for all $j = 2, \dots, k$. It follows from the arguments in the proof of Lemma 6.5 of [10] that given $\eta_j > 0$ there exists $C_j < \infty$

such that $\hat{\gamma}_j^{(n)}(x|_j) \leq C_j[(\hat{\gamma}_j(x|_j))^{1-\eta_j} \vee (\hat{\gamma}_j(x|_j))^{1+\eta_j}]$, and we replace $\vee$ by $+$, thus obtaining an upper bound. We thus obtain an upper bound for (6.21) in terms of 2^{k-1} sums of the form constant times

$$\sum_{x_1:\,\hat{\gamma}_1(x_1)\leq\delta_1} (\hat{\gamma}_1(x_1))^{1-\eta_1} \sum_{x_2} (\hat{\gamma}_2(x|_2))^{1\pm\eta_2} \cdots \sum_{x_k} (\hat{\gamma}_k(x|_k))^{1\pm\eta_k}. \tag{6.22}$$

Now, by choosing η_j small enough such that

$$\frac{\alpha_1}{1-\eta_1} < \frac{\alpha_2}{1\pm\eta_2} < \cdots < \frac{\alpha_k}{1\pm\eta_k} < 1, \tag{6.23}$$

one readily checks, e.g., by recoursing to Campbell's Theorem, that for every x_1

$$\sum_{x_2} (\hat{\gamma}_2(x|_2))^{1\pm\eta_2} \cdots \sum_{x_k} (\hat{\gamma}_k(x|_k))^{1\pm\eta_k} \tag{6.24}$$

is an $\frac{\alpha_2}{1\pm\eta_2}$-stable random variable, and finally that the random variable in (6.22), which is decreasing in δ_1, converges in probability to 0 as $\delta_1 \to 0$. We conclude it converges almost surely to 0 as $\delta_1 \to 0$, and (6.20) follows for the case where $i_1 = 2$ and $i_j \in \{1, 2\}$ for all $j = 2, \dots, k$.

Let us now analyse the expression in (6.21) when $\mathcal{L} := \{j = 1, \dots, k :\ i_j = 3\} \neq \emptyset$. It is argued in [10] (see discussion leading to (6.20) in that reference) that for $j \in \mathcal{L}$, $\hat{\gamma}_j^{(n)}(x|_j)$ is bounded above by constant times $c_j^{(n)}$. Let $k' = |\mathcal{L}|$ and let $i'_1 < \cdots < i'_{k'}$ be an enumeration of $\mathcal{L}$, and let $i''_1 < \cdots < i''_{k''}$ be an enumeration of $\{1, \dots, k\} \setminus \mathcal{L}$, $k'' = k - k'$. Then, arguing as above, (6.21) may be bounded above by a sum of $2^{k''}$ terms of the form

$$\prod_{j=1}^{k'} c_{i'_j}^{(n)} \sum_{x_{i'_1}=1}^{M_{i'_1}} \cdots \sum_{x_{i'_{k'}}=1}^{M_{i'_{k'}}} \left\{ \sum_{x_{i''_1}=1}^{M_{i''_1}} \left(\hat{\gamma}_{i''_1}^{(n)}(x|_{i''_1})\right)^{1\pm\eta_{i''_1}} \cdots \sum_{x_{i''_{k''}}=1}^{M_{i''_{k''}}} \left(\hat{\gamma}_{i''_{k''}}^{(n)}(x|_{i''_{k''}})\right)^{1\pm\eta_{i''_{k''}}} \right\}. \tag{6.25}$$

Again, choosing η's small enough, we have that the random variables within braces are iid $\frac{\alpha_{i''_1}}{1\pm\eta_{i''_1}}$-stable ones, and since the outer sums are over $\prod_{j=1}^{k'} M_{i'_j}$ terms, and, as one may readily check, $\prod_{j=1}^{k'} c_{i'_j}^{(n)} (\prod_{j=1}^{k'} M_{i'_j})^{(1\pm\eta_{i''_1})/\alpha_{i''_1}}$ decays polinomially in n to 0 as $n \to \infty$, by a standard argument, we have that the expression in (6.25) decays almost surely to 0 as $n \to \infty$, and (6.16) follows for general k by first taking $n \to \infty$ and then $\delta_1, \dots, \delta_k \to 0$.

It remains to check (5.7,5.8) as strong limits for the $\hat{\gamma}$ representations of the respective γ's. This is done in much the same way as for checking (5.6) above, so we will be rather sketchy. First note that the expressions in (5.7,5.8) can, after dividing the M's on the denominator inside the sum, and expanding the resulting products

$$\prod_{q=p+2}^{j} \left(\frac{1}{M_q} + \hat{\gamma}_{q-1}^{(n)}(x|_{q-1}) \right), \tag{6.26}$$

be both written as a sum over a fixed number of terms of the form

$$\sum_{x_1} \check{\gamma}_1^{(n)}(x_1) \cdots \sum_{x_\ell} \check{\gamma}_\ell^{(n)}(x|_\ell), \tag{6.27}$$

where $1 \leq \ell \leq k$ and $\check{\gamma}_j^{(n)}(x|_j)$ is either $\hat{\gamma}_j^{(n)}(x|_j)$ or $1/M_{j+1} = c_j^{(n)}$ for all $j = 1, \ldots, \ell$, with the latter case happening for at least one such j.

We can thus break each sum $\sum_{x_j}$ in three kinds as above (see (6.18)), with the superscript "(3)" applying also to the case where $\check{\gamma}_j^{(n)}(x|_j) = c_j^{(n)}$. The same arguments used above to estimate the latter cases of (6.21) (see paragraph of (6.25)) apply, since there is always a sum of the third kind, and the result follows.

□

Acknowledgements

LRF would like to thank the CMI, Université de Provence, Aix-Marseille I for hospitality and support during several visits in the last few years where this and related projects were developed. VG thanks NUMEC-USP for hospitality.

References

[1] Nieuwenhuizen, T.M.; Ernst, M.H. (1985) Excess noise in a hopping model for a resistor with quenched disorder, *J. Stat. Phys.* **41**, 773801

[2] Compte, A.; Bouchaud, J.-P. (1998) Localization in one-dimensional random random walks, *J. Phys. A: Math. Gen.* **31**, 61136121

[3] Fontes, L.R.G.; Isopi, M.; Newman, C. M. (2002) Random walks with strongly inhomogeneous rates and singular diffusions: convergence, localization and aging in one dimension, *Ann. Probab.* **30**, 579-604

[4] Ben Arous, G.; Černý, J.; Mountford, T. (2006) Aging in two-dimensional Bouchaud's model, *Probab. Theory Related Fields* **134**, 1–43

[5] Ben Arous, G.; Černý, J. (2007) Scaling limit for trap models on $\mathbb{Z}^d$, *Ann. Probab.* **35**, no. 6, 2356–2384.

[6] Barlow, M.; Černý, J. (2011) Convergence to fractional kinetics for random walks associated with unbounded conductances, *Probab. Theory Related Fields* **149**, 639–673.

[7] Bouchaud, J.-P.; Dean, D. S. (1995) Aging on Parisi's tree, *J. Phys. I France* **5**, 265–286

[8] Bovier, A.; Faggionato, A. (2005) Spectral characterisation of ageing: the REM-like trap model in the complete graph, *Ann. Appl. Probab.* **15**, 1997–2037

[9] Fontes, L.R.G.; Mathieu, P. (2008) K-processes, scaling limit and aging for the trap model on the complete graph, *Ann. Probab.* **36**, 1322-1358.

[10] Gayrard, V. (2010) Aging in reversible dynamics of disordered systems. I. Emergence of the arcsine law in Bouchaud's asymmetric trap model on the complete graph, arXiv:1008.3855v1 [math.PR]

[11] Bezerra, S.C.; Fontes, L.R.G.; Gava, R.J., Gayrard, V.; Mathieu, P. (2011) In preparation

[12] Ben Arous, G.; Bovier, A.; Gayrard, V. (2003) Glauber dynamics of the random energy model. I. Metastable motion on the extreme states, *Comm. Math. Phys.* **235**, no. 3, 379–425

[13] Ben Arous, G.; Bovier, A.; Gayrard, V. (2003) Glauber dynamics of the random energy model. II. Aging below the critical temperature, *Comm. Math. Phys.* **236**, no. 1, 1–54.

[14] Ben Arous, G.; Černý, J. (2008) The arcsine law as a universal aging scheme for trap models, *Comm. Pure Appl. Math.* **61**, 289–329

[15] Fontes, L.R.G.; Lima, P.H.S. (2009) Convergence of symmetric trap models in the hypercube, In: XVth International Congress on Mathematical Physics, 2006, Rio de Janeiro, *New Trends in Mathematical Physics*, Springer, 285-297

[16] Gayrard, V. (2010) Aging in reversible dynamics of disordered systems. II. Emergence of the arcsine law in the random hopping time dynamics of the REM, arXiv:1008.3849

[17] Ben Arous, G.; Gün, O. (2012) Universality and extremal aging for dynamics of spin glasses on subexponential time scales, *Comm. Pure Appl. Math.* **65**, 77127

[18] Ben Arous, G.; Bovier, A.; Černý, J. (2008) Universality of the REM for dynamics of mean-field spin glasses, *Comm. Math. Phys.* **282**, 663-695

[19] Bovier, A.; Gayrard, V. (2011) Convergence of clock processes in random environments and ageing in the p-spin SK model, to appear in *Ann. Probab.*

[20] Bovier, A.; Gayrard, V.; Svejda, A. (2011) Convergence to extremal processes in random environments and extremal ageing in the p-spin SK model, preprint

[21] Sasaki, M.; Nemoto, K. (2000) Analysis on Aging in the Generalized Random Energy Model, *J. Phys. Soc. Jpn.* **69**, 3045-3050

[22] Sasaki, M.; Nemoto, K. (2001) Numerical Study of Aging in the Generalized Random Energy Model, *J. Phys. Soc. Jpn.* **70**, 1099-1104

[23] Gayrard, V.; Gün, O. (2011) In preparation

[24] Ethier, S.N.; Kurtz, T.G. (1986) *Markov processes. Characterization and convergence*, Wiley

[25] Lepage, R.; Woodroofe, M.; Zinn, J. (1981) Convergence to a stable distribution via order statistics, *Ann. Probab.* **9**, 624632.